

# Time-Bin Qubits *via* Transformation of a Single Photon Field into Bunches of Pulses

R N SHAKHMURATOV<sup>1,2</sup>, F G VAGIZOV<sup>2</sup>, V A ANTONOV<sup>3</sup>, Y V RADEONYCHEV<sup>3</sup>, M O SCULLY<sup>4</sup>, AND O KOCHAROVSKAYA<sup>4</sup>

<sup>1</sup>*Kazan Physical-Technical Institute, Kazan, Russia*

<sup>2</sup>*Institute of Physics, Kazan Federal University, 18 Kremlyovskaya st., 420008, Kazan, Russia*

<sup>3</sup>*Institute of Applied Physics, Nizhny Novgorod, Russia*

<sup>4</sup>*Department of Physics and Astronomy, Texas A&M University, 77843-4242, College Station, TX, USA*

Contact Email: shakhmuratov@mail.ru

We propose a method to transform a single photon field into bunches of pulses with controllable timing and number of pulses in a bunch. The method is based on transmission of a photon through an optically thick single-line absorber vibrated with a frequency appreciably exceeding the width of the absorption line. The narrow spectrum of the incoming photon is “seen” by the vibrated absorber as a comb of equidistant spectral components separated by the vibration frequency. Tuning the absorber in resonance with  $m$ -th spectral component transforms the output radiation into bunches of pulses with  $m$  pulses in each bunch. We experimentally demonstrate the proposed technique with single 14.4 keV photons. This method opens a new way to the production of time-bin qubits. Proposals for construction of time-bin qubits with dimension higher than two by the pulse-bunching technique are discussed.