

**INTERNATIONAL CONFERENCE
DAYS ON DIFFRACTION 2014**

ABSTRACTS



May 26 – 30, 2014

St. Petersburg

ORGANIZING COMMITTEE

V. M. Babich /Chair/, A. S. Kirpichnikova /Secretary/,
T. V. Vinogradova /Visas/, N. V. Zalesskaya /Accommodation/,
I. V. Andronov, P. A. Belov, N. Ya. Kirpichnikova, A. P. Kiselev,
M. A. Lyalinov, O. V. Motygin, M. V. Perel, A. M. Samsonov,
V. P. Smyshlyaev, R. Stone, V. N. Troyan, N. Zhu

Conference e-mail: diffraction14@gmail.com

Web site: <http://www.pdmi.ras.ru/~dd/>

The conference is organized and sponsored by



St. Petersburg
Department of
V.A. Steklov
Institute of Mathematics



St. Petersburg State
University



The Euler International
Mathematical Institute



ITMO University



Russian Foundation
for Basic Research



IEEE Russia (Northwest)
Section AP/ED/MTT
Joint Chapter



Russian Academy of
Sciences

References

- [1] T. Z. Ismagilov, Second order scheme for Maxwell's equations with discontinuous electromagnetic properties. *Lect. Notes Comput. Sc.*, **7125**, 227–233 (2012).
- [2] T. Z. Ismagilov, A second-order scheme for Maxwell's equations with dielectric permittivity discontinuities and total field-scattered field boundaries. *Int. J. Comput. Math.*, **89**, 1378–1387 (2012).

An inverse eigenvalue problem of the theory of optical waveguides

Karchevskii E.M., Spiridonov A.O.

Kazan Federal University, Russia

e-mails: ekarchev@yandex.ru, sasha_ens@mail.ru

Inverse eigenvalue problems arise in a remarkable variety of applications, including system and control theory, geophysics, molecular spectroscopy, particle physics, structure analysis, and so on. An inverse eigenvalue problem concerns the reconstruction of a physical system from prescribed spectral data. The spectral data involved may consist of the complete or only partial information of eigenvalues or eigenvectors.

We present a new method for calculation of permittivity of dielectric materials using optical fiber's propagation constant measurements. This problem is an inverse eigenvalue problem of the optical waveguide theory. We obtain a mathematical model of eigenmodes of a weakly-guiding step-index arbitrarily shaped optical waveguide. By methods of the integral equations theory we prove that it is enough to measure the propagation constant of the fundamental eigenmode only at one frequency for the reconstruction of unknown dielectric constant of this waveguide. We present a new numerical algorithm for the calculation of the dielectric constant based on approximate solution of a nonlinear nonselfadjoint inverse eigenvalue problem for a system of weakly singular integral equations. The convergence and quality of this numerical method we prove by numerical experiments.

Monodromy of Heun equations with apparent singularities

Alexander Kazakov

St. Petersburg University of technology and design

St. Petersburg University of aerospace instrumentation

e-mail: a.kazakov@mail.ru

Heun and confluent Heun equations when one regular singularity is apparent singularity are under consideration. Corresponding connection matrices are calculated in explicit form.

The Dunkl–Darboux differential-difference operators and integrability

S. Khekalo

Moscow Regional State Institute of Humanities and Social Studies, 140410, Russian Federation, Moscow Region, Kolomna, Zelenaya str., 30

e-mail: khekalo@mail.ru

Let \mathfrak{F} be a space of the differentiable functions on \mathbb{R} ; \hat{s} be a reflection operator, so that $\hat{s}[f](x) = f(-x)$, $f(x) \in \mathfrak{F}$, and $\omega(x)$, $x \in \mathbb{R}$, be an even analytical function in an open domain $D_\omega \in \mathbb{R}$.

We consider the Dunkl–Darboux differential-difference operators in \mathbb{R}

$$\widehat{\nabla}_\omega = \frac{d}{dx} - (\ln |\omega(x)|)' \hat{s}. \quad (1)$$