



1128393

Global Experts Meeting on

Frontiers in Food Security & Nutrition

September 09-11, 2019 Edinburgh, Scotland

Enhancement of bone health using fish oil

*Amr S. Elbahnasawy^{*1,2}, E.R. Valeeva¹, Eman M. El-Sayed² and N.V. Stepanova¹*

¹ *Department of Bioecology, Hygiene and public health, Institute of Fundamental Medicine and Biology, Kazan Federal University, Kazan, Russia.*


² *Department of Nutrition and Food Sciences, National Research Centre, Dokki, Giza, Egypt.*

Abstract

Backgrounds: Osteoporosis is recognized by decreased bone density and deterioration in bone structure, causing a decrease in bone strength and an increase in bone fragility and fracture problems. Glucocorticoids (GCs) are considered as one of the main reasons of the secondary osteoporosis, and the resulting fractures cause significant morbidity. fish oil can decrease bone resorption, inhibit osteoclastogenesis, and prevent bone resorption. The aim of the current work was to evaluate the role of cod liver oil in preventing bone loss and osteoporosis in glucocorticoid-treated rats. **Methods:** The fatty acids profile of cod liver oil was analyzed, male rats were fed balanced diet and subdivided into 3 groups, 1- normal control, 2- prednisolone control, administrated prednisolone (10 mg/kg po daily); 3- fish oil, administrated prednisolone + cod liver oil (7% w/w). **Results:** Fatty Acid analyses showed that cod liver oil contained high levels of long chain ω -3 polyunsaturated fatty acids (PUFAs), eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA). Supplementation with cod liver oil helped to improve calcium levels in plasma, and suppress oxidative stress and inflammatory markers. Additionally, reduced bone resorption as reflected from the decreased levels of C-terminal telopeptide, and showed significant improvement in bone mineral density and normal histological results of bone cells compared to normal and prednisolone control. **Conclusions:** Our study indicated that feeding oil rich in PUFAs such as cod liver oil was able to modulate the potential effect of prednisolone in bone loss in rats. This may occur through some intracellular pathways, involving the improvement of calcium absorption, regulation of bone metabolism and the differentiation of the osteoblast and osteoclast, suppression of oxidative

stress and modulation of inflammatory response. Therefore, cod liver oil could be used as a natural approach to help in preventing bone loss associated with glucocorticoid therapy.

Keywords: Glucocorticoid-Induced Osteoporosis, cod liver oil, ω -3 polyunsaturated fatty acids.



James Walker

Program Ambassador | Food Security 2019, Scotland
+44 203 455 4929, foodsecurity@frontiermeetings.com

Organizing Committee Members: Food Security 2019, Scotland

Dr. Catherine Keske

University of California, USA

Dr. Yoshie Sano

Washington State University Vancouver, Canada

Dr. Gilbert Onionkiton Adjimoti

University of abomey-calavi, France

Dr. Jean Pierre Poulain

University of Toulouse, France

16 Medway Road | Crayford, Dartford | DA1 4PN | London, UK Ph.: +44 203 455 4929
E: info@frontiersmeetings.com | W: <https://frontiersmeetings.com/conferences/foodsecurity/>