56TH ANNUAL SCIENTIFIC MEETING – 8–10 June 2022, Bari, Italy



Cholangiocarcinoma - 9 faces of the killer It shows cholangiocarcinoma, an aggressive bile duct tumour with dismal prognosis, It was captured during magnetic resonance cholangiopancreatography (MRCP) Piotr Milkiewicz, Warsaw Poland





THE JOURNAL OF THE EUROPEAN SOCIETY FOR CLINICAL INVESTIGATION

EDITOR-IN-CHIEF	Hendrik M. Nathoe	EDITORIAL ADDRESS Bolognalaan 34 3584 CJ Utrecht, The Netherlands	
DEPUTY EDITOR-IN-CHIEF	Fabrizio Montecucco		
MANAGING EDITORS	Ingrid Meijer. Utrecht, The Netherlands Debbie Barneveld Binkhuijsen. Utrecht, The Netherlands	Email: EJCIOffice@esci.eu.com	
PUBLISHER'S OFFICE	Production Editor: Arthi Riya eci@wiley.com		

Aims & Scope

The European Journal of Clinical Investigation (EJCI), in publication since 1970, is a peer-reviewed general-interest biomedical journal with a broad readership. It is the official journal of the European Society for Clinical Investigation (ESCI) and it is published monthly by Wiley. It considers any original contribution from the most sophisticated basic molecular sciences to applied clinical and translational research and evidence-based medicine across a broad range of subspecialties. The EJCI publishes reports of high-quality research that pertain to the genetic, molecular, cellular, or physiological basis of human biology and disease, as well as research that addresses prevalence, diagnosis, course, treatment, and prevention of disease. We are primarily interested in studies directly pertinent to humans, but submission of robust in vitro and animal work is also encouraged. Interdisciplinary work and research using innovative methods and combinations of laboratory, clinical, and epidemiological methodologies and techniques is of great interest to the journal. Several categories of manuscripts (for detailed description see below) are considered: editorials, original articles (also including randomized clinical trials, systematic reviews and meta-analyses), reviews (narrative reviews), opinion articles (including debates, perspectives and commentaries); and letters to the Editor.

Copyright and Copying

Copyright © 2022 Stichting European Society for Clinical Investigation Journal Foundation. Published by John Wiley & Sons Ltd. All rights reserved. No part of this publication (apart from articles marked 'OnlineOpen') may be reproduced, stored or transmitted in any form or by any means without the prior permission in writing from the copyright holder. Authorization to copy items for internal and personal use is granted by the copyright holder for libraries and other users registered with their local Reproduction Rights Organisation (RRO), e.g. Copyright Clearance Center (CCC), 222 Rosewood Drive, Danvers, MA 01923, USA (http://www.copyright.com), provided the appropriate fee is paid directly to the RRO. This consent does not extend to other kinds of copying such as copying for general distribution, for advertising and promotional purposes, for creating new collective works or for resale. Special requests should be addressed to: permissionsuk@wiley.com.

Disclaimer

The Publisher, European Society for Clinical Investigation and Editors cannot be held responsible for errors or any consequences arising from the use of information contained in this journal; the views and opinions expressed do not necessarily reflect those of the Publisher, European Society for Clinical Investigation and Editors, neither does the publication of advertisements constitute any endorsement by the Publisher, European Society for Clinical Investigation and Editors of the products advertised.

ISSN 0014-2972 (Print) ISSN 1365–2362 (Online)

THE JOURNAL OF THE EUROPEAN SOCIETY FOR CLINICAL INVESTIGATION

EDITORIAL BOARD

Editor-in-Chief Hendrik Nathoe Utrecht, The Netherlands

Deputy Editor-in-Chief Fabrizio Montecucco Genoa, Italy

Managing Editors Ingrid Meijer Utrecht, The Netherlands Debbie Barneveld Binkhuijsen Utrecht, The Netherlands

Deputy Editors Federico Carbone Genoa, Italy Yvonne Döring Bern, Switzerland Baris Gencer Boston, USA Luca Liberale Zurich, Switzerland Vilma Sardão Coimbra, Portugal

Section Editors

Johann Auer Braunau, Austria Leonilde Bonfrate Bari, Italy Marco Canepa Genoa, Italy Agostino Di Ciauli Bari, Italy Leo Koenderman Utrecht, The Netherlands Igor Koncar Belgrade, Serbia Marcin Krawczyk Warsaw, Poland Matteo Lambertini Genoa, Italy Paulo Oliveira Coimbra, Portugal Piero Portincasa Bari, Italy Voahanginirina Randriamboavonjy Frankfurt, Germany Ruslan Rust Zurich, Switzerland Thomas Schindler St Louis, USA Angela Angela Sciacqua Catanzaro, Italy Vanessa Smith Ghent, Belgium Markus Sperandio Munich, Germany Mario Strazzabosco New Haven, USA Marcel Twickler Antwerp, Belgium Gemma Vilahur Barcelona, Spain José Viña Valencia, Spain Hiroshi Yoshida Chiba, Japan Carmine Zoccali Reggio Calabria, Italy

European Society for Clinical Investigation



Founded 12 February 1967

Officers and members of the ESCI Council 2020

PRESIDENT

Paulo Oliveira, Coimbra, Portugal

PAST PRESIDENT Hendrik Nathoe, Utrecht, The Netherlands

VICE-PRESIDENT Voahangy Randriamboavongy, Frankfurt, Germany

SECRETARY-TREASURER Jakub Regieli, The Netherlands

COUNCILLORS

Federico Carbone, Genoa, Italy Marcin Krawczyk, Homburg, Germany Nahid Tabibzadeh, Paris, France Gemma Vilahur, Barcelona, Spain

ESCI CENTRAL OFFICE

Marliese Greven Debbie Barneveld Binkhuijsen Utrecht, The Netherlands esci@umcutrecht.nl

Objectives

The objectives of the Society are the advancement of medical practice through science; the cultivation of clinical research by the methods of the natural sciences; the correlation of science with the art of medical practice; the fostering of high standards of ethical practice and investigation; and the diffusion of a spirit of fraternity and international co-operation among and through its members.

Membership

Any person who has been actively involved in medical research in Europe, for example by involvement in original investigations in clinical or allied sciences of medicine, can apply for membership of ESCI. Members will receive the *EJCI* monthly. Any company, corporation, organisation, or trust can apply for corporate membership in the Society.

Meetings

The Society holds one Annual Scientific Meeting each year, usually in April/May. Meetings are held in different cities in Europe. Other Society's activities include the sponsorship of Workshops and Postgraduate Courses and the encouragement of the exchange of Scientists between Laboratories.

Contact information ESCI

Enquiries regarding membership of the Society or any of its activities should be addressed to: European Society for Clinical Investigation (ESCI), Central Office, Bolognalaan 34, 3584 CJ Utrecht, Netherlands. Tel: +31 88 755 8787 esci@umcutrecht.nl; www.esci.eu.com

Volume 52

Supplement 1

June 2022

56TH ANNUAL SCIENTIFIC MEETING – 8–10 June 2022, Bari, Italy

These abstracts have been published as they were received via online electronic submission. Every effort has been made to reproduce faithfully the abstracts as submitted. However, no responsibility is assumed by the organizers for any injury and/or damage to persons or property as a matter of products liability, negligence or otherwise, or from any use or operation of methods, products, instructions or ideas contained in the materials herein. Because of the rapid advances in medical sciences, we recommend that independent verification of diagnoses and drug doses should be made.

Contents

		Page number	
ABSTRACTS			
S1	The Gut-Liver Body Axis & Lifestyles: Lessons from Frontline Biomedical Research	8	
S2	Mitochondria	34	
S3	Cardiovascular & Metabolic Diseases	62	
S4	Bioinformatics and Computational Biology for Biomedicine	88	
S5	Clinical Ultrasonography: Tips & Tricks	103	
S6	Membrane Channels and Transporters: Translating Basic Research to New Drug Discovery		
	and Preclinical Development	106	
S7	Regenerative - Genomic Medicines	124	
S8	Multidisciplinary - Collaborative Clinical Investigation Between Medicine & Surgery	139	
S9	Transitional, Translational Aspects and Genetics of FMF	159	
S10	Microbiome, Metabolome and Lifestyles: More to Know	163	
S11	Focus on Gender Medicine	171	

THE JOURNAL OF THE EUROPEAN SOCIETY FOR CLINICAL INVESTIGATION

EDITORIAL BOARD

Editor-in-Chief Hendrik Nathoe Utrecht, The Netherlands

Deputy Editor-in-Chief Fabrizio Montecucco Genoa, Italy

Managing Editors Ingrid Meijer Utrecht, The Netherlands Debbie Barneveld Binkhuijsen Utrecht, The Netherlands

Deputy Editors

Federico Carbone Genoa, Italy Yvonne Döring Bern, Switzerland Baris Gencer Boston, USA Luca Liberale Zurich, Switzerland Vilma Sardão Coimbra, Portugal

Section Editors

Johann Auer Braunau, Austria Leonilde Bonfrate Bari, Italy Marco Canepa Genoa, Italy Agostino Di Ciauli Bari, Italy Leo Koenderman Utrecht, The Netherlands Igor Koncar Belgrade, Serbia Marcin Krawczyk Warsaw, Poland Matteo Lambertini Genoa, Italy Paulo Oliveira Coimbra, Portugal Piero Portincasa Bari, Italy Voahanginirina Randriamboavonjy Frankfurt, Germany Ruslan Rust Zurich, Switzerland Thomas Schindler St Louis, USA Angela Angela Sciacqua Catanzaro, Italy Vanessa Smith Ghent, Belgium Markus Sperandio Munich, Germany Mario Strazzabosco New Haven, USA Marcel Twickler Antwerp, Belgium Gemma Vilahur Barcelona, Spain José Viña Valencia, Spain Hiroshi Yoshida Chiba, Japan Carmine Zoccali Reggio Calabria, Italy

cases. Decrease in myocardial contractility compared to the initial values was not observed during the entire reperfusion period (LVDPinitial = 52 ± 5.2 mmHg and LVDPreperfusion = 58 ± 5.8 mmHg), what allows us to conclude about the effectiveness of myocardial protection by the new CPS.

Conclusions: In our experiment on a model of an isolated rat heart, which is widely used for the study of various CPR, we showed that the new solution is able to quickly and effectively cause myocardial plegia, and also does not interfere with the rapid and full recovery of its function after the start of reperfusion. This paper has been supported by the Kazan Federal University Strategic Academic Leadership Program (PRIORITY-2030)

56ASM-0099 | The effect of α2-adrenoreceptors on the isolated rats heart contractility with a model of myocardial infarction in the acute stage

<u>L. Khisamieva;</u> A. Kuptsova; R. Bugrov; N. Ziyatdinova; T. Zefirov

Kazan Federal University, Department of Human Health Protection, Kazan, Russia C.I.S.

Background: Stimulation of α 2-adrenergic receptors (α 2-AR) has a direct cardioprotective effect. The researchers revealed an increase in the expression of α 2-AR subtypes in rats with spontaneous hypertension, as well as the dys-function of α 2-AR and the inefficiency of signaling pathways associated with α 2-AR. The aim of this study was to investigate the effect of α 2-AR stimulation of the isolated heart contractility with a myocardial infarction (MI) model 24 hours after the influence.

Materials and Methods: Ex vivo experiments were performed on isolated hearts of intact rats (n = 7) and rats with a model of MI (n = 10) 24 hours after the operation. IM is reproduced by ligation of the anterior branch of the left coronary artery. Agonist α_2 -AR clonidine hydrochloride was used in concentration of 10⁻⁹ Mol. The data were statistically processed using Student's t-test.

Results: A comparative analysis of a_2 -AR stimulation revealed that in intact rats, the agonist reduced contractility by 21% (p < 0.05), and with the MI model, 24 hours after modeling, it increased by 38% (p < 0.01).

Conclusions: Thus, in our experiments α 2-AR stimulation revealed multidirectional effects on inotropy reaction of intact rats isolated heart and rats with the MI model. The study was supported by Russian Science Foundation (grant No. 21-15-00121, https://rscf.ru/project/21-15-00121/)

56ASM-0100 | If-currents are involved in heart contractility regulation of rats with acute myocardial infarction model

L. Khisamieva; R. Bugrov; A. Kuptsova; N. Ziyatdinova; T. Zefirov

Kazan Federal University, Department of Human Health Protection, Kazan, Russia C.I.S.

Background: Many factors have been identified in the development of coronary heart disease, such as hypertension, diabetes, smoking and an increase in heart rate. "Funny currents" (If-currents) play a key role in the electrogenesis of cardiomyocytes. The density of If currents increase in atrial and ventricular myocytes in patients with heart failure. The aim of this study was to investigate the effect of the blockade If-currents on the contractility of isolated heart with a model of myocardial infarction (MI) 24 hours after experimental influence.

Materials and Methods: Ex vivo experiments were performed on Langendorff-isolated hearts of intact (n = 7), sham-operated (n = 6) rats and with a model of MI (n = 10) 24 hours after experimental influence. IM is reproduced by ligation of the anterior branch of the left coronary artery. To block If currents, ZD7288 was used at a concentration of 10^{-9} Mol. The data were statistically processed using Student's t-test.

Results: Blockade of If-current revealed that in intact rats group ZD7288 increased contractility by 47% (p < 0.05), in sham-operated animals – by 23% (p < 0.01), in the MI model 24 hours after the simulation increased by 20% (p < 0.01).

Conclusions: Thus, our study showed that in the group of rats with a model of acute myocardial infarction, the effect of If on the contractility of the isolated heart decreases. The study was supported by Russian Science Foundation (grant No. 21-15-00121, https://rscf.ru/ project/21-15-00121/)

56ASM-0101 | Analysis of rats myocardial contraction force at different stages of experimental infarction

<u>L. Khisamieva</u>; R. Bugrov; A. Kuptsova; N. Ziyatdinova; T. Zefirov

Kazan Federal University, Department of Human Health Protection, Kazan, Russia C.I.S.

Background: The study of experimental models of myocardial infarction (MI) is necessary for the development of innovative methods of treatment of this disease. The aim of this study is to conduct a comparative analysis of the

WILEY-

effect of myocardial infarction on the contractility of the isolated rat heart at different stages of MI.

Materials and Methods: The experiments were performed ex vivo on isolated hearts of intact and rats with a model of MI after 1 day, 54 days and 120 days after the simulation. MI was performed according to the classical technique - ligation of the anterior branch of the left coronary artery. The contractile activity was studied on the Langendorff System. The data were statistically processed using Student's t-test.

Results: A comparative analysis of the effect of MI on the initial values of contraction force in the studied groups revealed that in rats the contraction force decreased one day after MI and tended to increase 54 and 120 days after the simulation of MI.

Conclusions: Thus, it was shown that at different stages of the postinfarction period, multidirectional changes of the isolated rat heart myocardium contractions force are observed. The study was supported by Russian Science Foundation (grant No. 21-15-00121, https://rscf.ru/project/21-15-00121/)

56ASM-0102 | Effect of HCN channel blocker in the regulation of chronotropic effects in rats with limited motor activity

<u>M. Sungatullina;</u> R. Zaripova; R. Shakirov; N. Ziyatdinova; T. Zefirov Kazan Federal University, Department of Human Health Protection, Kazan, Russia C.I.S.

Background: In the modern world, limitation of motor activity is an acute problem, because there are many reasons leading to this way of life. Hypokinesia causes atrophy of the musculoskeletal system, complicates the digestive, respiratory and cardiovascular systems. The involvement of HCN channels in the mechanism of heart rhythm acceleration has been shown. It is interesting to see the effect of blockade of If-currents and their role in the regulation of chronotropy of the heart against the background of increased heart rate (HR) in response to hypokinesia.

Materials and Methods: The experiments were conducted on two groups of rats: 1- control group, rats 7 weeks old; 2 - experimental group, rats with restriction of motor activity for 30 days. This effect was achieved by placing 3-week-old rats in penal cages under conditions of increasing hypokinesia. The effect of If blocker ZD7288 (10⁻⁹M and 10⁻⁶M) on chronotropic effects was studied using Langendorff PowerLab8/35 (ADInstruments, Australia). **Results:** After the introduction into the perfused solution ZD7288 (10⁻⁹M), a decrease in heart rate by 15% was observed in control rats (p < 0.01) and by 11% (p < 0.05) in the experimental group. The blocker If in concentrations 10⁻⁶M decreased heart rate in the control group by 28% (p < 0.01) and by 17% in experimental group.

Conclusions: If-current blocker ZD7288 at all concentrations caused a decrease in heart rate in control rats and rats, with limitation of motor activity. However, more pronounced changes in heart rate were observed in the control group of rats and after application of the maximum concentration. It is possible that in rats with limited motor activity, against the background of an increase in heart rate, the density of HCN channels decreases compensatory, which leads to decrease their role in the regulation of heart rate. This paper has been supported by the Kazan Federal University Strategic Academic Leadership Program (PRIORITY-2030)

56ASM-0103 | Isolated rat heart after restriction of motor activity and recovery

<u>M. Sungatullina;</u> R. Zaripova; L. Mosolov; N. Ziyatdinova; T. Zefirov *Kazan Federal University, Department of Human Health Protection, Kazan, Russia C.I.S.*

Background: Restriction of motor activity becomes a medical and social problem. Restriction of muscle activity leads to violations of all organ systems of the human body. Namely, in the cardiovascular system, prolonged restriction of motor activity leads to coronary vessels, the heart muscle weakening, decreasing of the heart energy potential and minute volume. The aim of our study was to study possible age-related changes in the parameters of the isolated rat heart after hypokinesia and subsequent recovery. Materials and Methods: Restriction of motor activity was carried out by placing animals in pencil cases in conditions of increasing hypokinesia for 30 days. The recovery stage after hypokinesia for 14 days was carried out in order to study the mechanisms of adaptation of the animal to changes in the motor regime. The following parameters of the isolated heart were recorded - the pressure developed in the left ventricle (LVL), heart rate (HR) and coronary flow (CP) on the Langendorff PowerLab 8/35 unit (ADInstruments, Australia). Statistical processing was carried out in Excel, the reliability was determined using the Student's t-test.

Results: After hypokinesia, unidirectional changes in the parameters of the isolated heart were observed in 7-week-old and adult rats: a decrease in the parameters of the LVL, CP and an increase in heart rate. However,