




KAZAN FEDERAL UNIVERSITY

Russian Gravitational Society
5-100 Russian Academic Excellence Project



**4-я Международная зимняя школа-семинар
по гравитации, космологии и астрофизике
«Петровские чтения-2018»
Программа и тезисы докладов**

**4th International Winter School-Seminar
on gravity, cosmology, and astrophysics
«Petrov School-2018»
Program and Abstracts**

Казань, 2018

4-я Международная зимняя школа-семинар по гравитации, космологии и астрофизике «Петровские чтения-2018». Программа и тезисы докладов международной научной школы-семинара. — Казань: Изд-во КФУ, 2018. — 64 с.

Сборник содержит программу и тезисы докладов участников 4-ей Международной зимней школы-семинара по гравитации, космологии и астрофизике «Петровские чтения-2018». В материалах представлены работы ведущих специалистов и начинающих исследователей из научных центров России, ближнего и дальнего зарубежья, посвященные исследованиям в ОТО и модифицированных теориях гравитации, теоретической и наблюдательной космологии, релятивистской астрофизике. Международная зимняя школа-семинар проходила в Казанском университете с 26 ноября по 1 декабря 2018 года.

Сборник адресован научным работникам, аспирантам и молодым ученым, специализирующимся в области теории гравитации, космологии и астрофизики, а также для студентов старших курсов естественнонаучных направлений.

Школа-семинар «Петровские чтения-2018» проводится за счет средств субсидии, выделенной в рамках государственной поддержки Казанского (Приволжского) федерального университета в целях повышения его конкурентоспособности среди ведущих мировых научно-образовательных центров. Проведение «Петровских чтений-2018» нацелено на реализацию научно-образовательных задач в рамках стратегической академической единицы «Астровывзов».

СОДЕРЖАНИЕ

ПРОГРАММА / PROGRAM

7

ЛЕКЦИИ ШКОЛЫ / LECTURES

17

<i>A.A. Starobinsky.</i> Search for quantum-gravitational effects in the early and present Universe	17
<i>S. Capozziello.</i> Dark Energy and Dark Matter as Curvature Effects	17
<i>I.I. Tkachev.</i> Dark matter models. From graviton to WIMPZILLA	18
<i>T. Prokopec.</i> Quantum effects in cosmological perturbation theory	18
<i>S.D. Odintsov.</i> Unifying inflation with dark energy: the case of modified gravity . .	18
<i>M.S. Volkov.</i> New theories with massive gravitons and new Horndeski-type theories	19
<i>Eu. Babichev.</i> Modern scalar-tensor theories of gravity	19
<i>A. Vikman.</i> Dark Energy and Dark Matter as Lagrange Multipliers	20
<i>V. Berezhin.</i> Higher derivative gravity: Alexey Starobinsky and beyond	20
<i>K.A. Bronnikov.</i> Nonlinear electrodynamics, black holes and wormholes	22
<i>I.L. Buchbinder.</i> Renormalization of quantum field theory in curved space-time. Brief lecture course for beginners	23
<i>D.V. Gal'tsov.</i> Disformal dualities and the problem of spacetime singularities	23
<i>G.L. Klimchitskaya, V.M. Mostepanenko.</i> Axion-like particles in gravitation, astrophysics and cosmology	24
<i>D.S. Gorbunov.</i> Dark Matter: thermal and non-thermal	24

ТЕЗИСЫ ДОКЛАДОВ / ABSTRACTS

26

<i>S. Alexeyev, B. Latosh, V. Echeistov.</i> Starobinsky Model with Vanishing Cosmological Constant at Galaxy Clusters Scales	26
<i>T.Yu. Alpin.</i> PP-wave modes in an electromagnetically active dynamic aether: Birefringence and anomalies of the phase velocity	26
<i>S.N. Andrianov, R.A. Daishev, S.M. Kozyrev, B.P. Pavlov.</i> Geometrization of stress-energy tensor	27
<i>A.V. Aminova, D.R. Khakimov.</i> On five-dimensional spaces with projective symmetries	28
<i>N.A. Avdeev, P.I. Dyadina.</i> Testing of hybrid metric-Palatini $f(R)$ -gravity in eccentric binary pulsars	28
<i>K. Boshkayev.</i> A model for a dark matter core at the galactic center	28
<i>S.V. Chervon.</i> Chiral cosmological models and modified theories of gravitation. . .	29
<i>A.O. Efremova.</i> Interaction between spinor field and dynamic aether	30
<i>J.C. Fabris, G. Brando, F.T. Falciano, O. Galkina.</i> Stiff Matter Solution in Brans-Dicke Theory and The General Relativity Limit	31
<i>I.V. Fomin.</i> GR-like inflationary models in modified gravity theories	31
<i>A.A. Galimova.</i> Dilatonic extension of the Einstein-Aether theory	32
<i>D.V. Gal'tsov, S.M. Zhidkova.</i> Isolating singularities in Palatini version of kinetically coupled scalar-tensor theory	32

<i>D.E. Groshev, A.B. Balakin.</i> Polarization and stratification of axionically active plasma in a dyon magnetosphere	33
<i>A. Gusev.</i> Chandler wobble and free core nutation of seven single pulsars	33
<i>Yu. Ignat'ev.</i> The Peculiarities of the Cosmological Models Based on Nonlinear Classical and Phantom Fields	35
<i>V.D. Ivashchuk, A.A. Kobtsev.</i> Exact solutions in multidimensional Einstein-Gauss-Bonnet model with cosmological term	36
<i>R.N. Izmailov, K.K. Nandi.</i> Can massless wormholes mimic Schwarzschild black hole in the strong field lensing?	36
<i>A. Kamaletdinov, E. Esipova.</i> Investigation of possible cosmological constraints of SIMP-like dark matter models	37
<i>R.Kh. Karimov, R.N. Izmailov.</i> Accretion disks properties of charged black holes	38
<i>R.L. Khazhin.</i> Memory effects in a magnetized cosmic dark fluid	39
<i>A.A. Kirillov, E.P. Savelova.</i> Wormhole as a natural accelerator of high energy cosmic rays	39
<i>A.A. Kirillov, E.P. Savelova.</i> Virtual wormholes and dynamical Pauli-Villars regularization	40
<i>A.S. Koshelev.</i> UV-completion of the Starobinsky inflation in the framework of analytic infinite derivative gravity	41
<i>K.K. Nandi, R.N. Izmailov.</i> Stability of Ellis-Bronnikov wormhole: Is it observer-location dependent?	41
<i>A.A. Popov, D.S. Lisenkov.</i> Vacuum polarization of a quantized scalar field in the thermal state in a long throat	42
<i>E.O. Pozdeeva.</i> Multifield inflation	44
<i>B. Saha.</i> Spinor fields in spherically symmetric space-time.	44
<i>A.F. Shakirzyanov.</i> Einstein-aether-axion theory: How does dynamic aether regulate the state of axionic dark matter?	44
<i>M.A. Skugoreva, A.V. Toporensky.</i> On Kasner solution in Bianchi I $f(T)$ cosmology	45
<i>I. Timiryasov, Sh. Ejima, M. Shaposhnikov.</i> Parameter space of baryogenesis in the ν MSM	45
<i>A. Tokareva, D. Gorbunov.</i> R^2 gravity makes Higgs inflation consistent	46
<i>A.V. Toporensky, O. Zaslavsky.</i> The 'river model' of black hole and its applications	46
<i>A. Urazalina, V. Dzhunushaliev.</i> Regular solutions in GR with potentials of the 4th, 6th and 8th orders	46
<i>S.Yu. Vernov.</i> Cosmological models with the Gauss-Bonnet term	47
<i>A.V. Yurov, V.A. Yurov.</i> Quantum Cosmology without a Wave function	48
<i>F. Zaripov.</i> Oscillating solutions of a modified theory of induced gravity in cosmology and in centrally symmetric metric	48
<i>Д.Л. Аббаров.</i> Дзета-модель волчка Джанибекова и его аналогия с гравитоном	49
<i>А.В. Аминова, А.И. Гарафутдинова, М.Х. Люлинский.</i> Решение полевых уравнений для суперпространства Минковского	50
<i>К.А. Большакова, С.В. Червон.</i> Инфляционные решения с заданными потенциалами в тензорно-мульти-скалярной теории гравитации.	50

<i>В.О. Гладышев, В.И. Пустовойт, В.С. Горелик, В.Л. Кауц, А.В. Каютенко, Е.А. Шарандин.</i> О генерации и детектировании высокочастотных гравитационных волн в диэлектрических средах при их возбуждении оптическим излучением	51
<i>В.М. Журавлев.</i> Принцип материальности пространства и фундаментальные поля	52
<i>С.В. Зуев.</i> Вероятностный численный метод решения ОДУ с применением квантовых вычислений	54
<i>А.С. Ильин.</i> Dark energy and dark matter interaction: kernels of Volterra type and coincidence problem	55
<i>Р.Ф. Лукманова.</i> Характеристики линзирования безмассовыми кротовыми норами в пределе слабого поля	55
<i>И.С. Нургалиев.</i> Несингулярная космология и ее уроки.	56
<i>В.Ф. Панов, О.В. Сандакова, Е.В. Кувшинова, Д.М. Янишевский, М.Р. Черемных.</i> Космология с вращением	57
<i>В.А. Прокопов, С.О. Алексеев.</i> Приливной заряд и поправки к нему в астрофизике чёрных дыр	58
<i>Э.А. Тагиров.</i> Принцип Гамильтона vs общая ковариантность при квантовании натуральных гамильтоновых систем	58
<i>П.В. Третьяков.</i> Spin connection and cosmological perturbations in teleparallel gravity.	59
<i>С.В. Червон, А.А. Фатахов.</i> Киральные космологические модели с неминимальным взаимодействием	59
<i>С.В. Червон, И.В. Фомин, Т.И. Майорова.</i> Космологические решения $f(R)$ гравитации с высшими производными	60
АВТОРСКИЙ УКАЗАТЕЛЬ	62

medium version of the Pound-Rebka experiment in gravity. This parallel lends validity to the "effective medium" approach. The probability of stability of a certain object is redefined in terms of reflection (R) and transmission (T) coefficients of the perturbations hitting the throat or horizon. Our conclusion is that there is a non-zero probability that EB and phantom wormholes could appear stable or unstable depending on the *location of observers*, leading to the possibility of *ghost wormholes* (similar to ghost stars). Schwarzschild horizon, however, would always appear stable ($R = 1$, $T = 0$) to observers independently of their locations. Phantom wormholes of bounded mass in the extreme limit $a \rightarrow -1$ are also shown to be stable just as the Schwarzschild black hole is. We shall propose a thought experiment showing that our non-deterministic results could be numerically translated into observable deterministic signatures of ghost wormholes.

References

1. H.G. Ellis, J. Math. Phys. **14**, 104 (1973); **15**, 520(E) (1974).
2. K.A. Bronnikov, Acta Phys. Polon. B **4**, 251 (1973).
3. J.A. González, F.S. Guzmán and O. Sarbach, Class. Quant. Grav. **26**, 015010 (2009).
4. J.A. González, F.S. Guzmán and O. Sarbach, Class. Quant. Grav. **26**, 015011 (2009).
5. F.S. N. Lobo, F. Parsaei and N. Riazi, Phys. Rev. D **87**, 084030 (2013).
6. S.V. Sushkov, Phys. Rev. D **71**, 043520 (2005).
7. K. K. Nandi, A. A. Potapov, R. N. Izmailov, A. Tamang, and J. C. Evans, Phys. Rev. D **93**, 104044 (2016).

VACUUM POLARIZATION OF A QUANTIZED SCALAR FIELD IN THE THERMAL STATE IN A LONG THROAT

A.A. Popov¹, D.S. Lisenkov²

¹ *apopov@kpfu.ru*; Institute of Mathematics and Mechanics, Kazan Federal University, Kazan, 420008, Russia

² *lesman1985@gmail.com*; Institute of Mathematics and Mechanics, Kazan Federal University, Kazan, 420008, Russia

The study of vacuum polarization effects in strong gravitational fields is a pertinent issue since such effects may play a role in the cosmological scenario and the construction of a self-consistent model of black hole evaporation. These effects may be taken into account by solving the semiclassical backreaction equations,

$$G_{\nu}^{\mu} = 8\pi \langle T_{\nu}^{\mu} \rangle, \quad (1)$$

where $\langle T_{\nu}^{\mu} \rangle$ is the expectation value of the stress-energy tensor operator for the quantized fields.

The main difficulty in the theory of semiclassical gravity is that the effects of the quantized gravitational field are ignored. The popular solution for this problem is to justify ignoring the gravitational contribution by working in the limit of a large number of fields,

in which the gravitational contribution is negligible. Another problem is that the vacuum polarization effects are determined by the topological and geometrical properties of spacetime as a whole or by the choice of quantum state in which the expectation values are taken. It means that the calculation of the functional dependence of $\langle T_{\nu}^{\mu} \rangle_{ren}$ on the metric tensor in an arbitrary spacetime presents formidable difficulty. Only in some spacetimes with high degrees of symmetry for the conformally invariant fields $\langle T_{\mu\nu} \rangle_{ren}$ can be computed and equations (1) can be solved exactly [1–5]. Let us stress that the single parameter of length dimensionality in such a problem is the Planck length l_{pl} . This implies that the characteristic scale l of the spacetime curvature (which correspond to the solution of equations (1)) can differ from l_{pl} only if there is a large dimensionless parameter. As an example of such a parameter one can consider a number of fields, the polarization of which is a source of spacetime curvature¹. For the quantized scalar field such a parameter can be the coupling constant ξ of scalar field to the curvature of spacetime. In the case of $|\xi| \gg 1, \xi < 0$ the vacuum fluctuations of a quantized scalar field can determine the curvature of spacetime so-called *a long throat* [6].

In this work we have shown that in a long throat the effect of vacuum polarization of a quantized scalar field in the thermal state does not depend on temperature and conditions at infinity. This implies that in considered situation $\langle \varphi^2 \rangle$ is a local quantity for any finite mass m of the quantized field, including $m = 0$.

References

1. Starobinsky A. A. *A new type of isotropic cosmological models without singularity* // Phys. Lett. B – 1980. – Vol. 91. – Is. 1. – p. 99-102.
2. Mamaev S. G., Mostepanenko V. M. *Isotropic cosmological models determined by vacuum quantum effects* // Sov. Phys.–JETP 1980. – Vol. 51 (1). – p. 9-13.
3. L. A. Kofman, V. Sakhni, and A. A. Starobinskii *Anisotropic cosmological model created by quantum polarization of vacuum* // Sov. Phys.–JETP 1980. – Vol. 58 (6). – p. 1090.
4. Kofman L. A., Sahni V. *Hew self-consistent solution of the Einstein equations with one-loop quantum gravitational corrections* // Phys. Lett. B – 1983. – Vol. 127. – p. 197-200.
5. Kofman L. A., Sahni V. *Some self-consistent solutions of the Einstein equations with one-loop quantum gravitational corrections: $G_{ik} = 8\pi G \langle T_{ik} \rangle_{vac}$* // Phys. Lett. A – 1986. – Vol. 117. – p. 275-278.
6. Popov A. A. *Long throat of a wormhole created from vacuum fluctuations* // Classical and Quantum Gravity – 2005. – Vol. 22. – № 24. – p. 5223-5230.

¹ Here and below it is assumed, of course, that the characteristic scale of change of the background gravitational field is sufficiently greater than l_{pl} so that the very notion of a classical spacetime still has some meaning.