

Kazan Federal University

Zavoiskii Physical-Technical Institute, FRC Kazan Scientific Center  
of RAS

International Conference "Magnetic Resonance - Current State and  
Future Perspectives" and satellite XXI International Youth  
Scientific School "Actual problems of magnetic resonance and its  
application"

devoted to the 75-th anniversary of the discovery of Electron Paramagnetic  
Resonance by E.K. Zavoiskii

Book of  
**ABSTRACTS**

**September 23-27, 2019 Kazan, Russia**

## An Investigation of the Conformational Properties of Mefenamic Acid in DMSO by Two-dimensional NMR

I.A. Khodov<sup>1,2</sup>, S.V. Efimov<sup>2</sup>, K.V. Belov<sup>1</sup>, L.A.E. Batista de Carvalho<sup>3</sup>

<sup>1</sup>G.A. Krestov Institute of Solution Chemistry of the Russian Academy of Sciences, Ivanovo, Russian Federation

<sup>2</sup>Institute of Physics, Kazan Federal University, Kazan, Russian Federation

<sup>3</sup>University of Coimbra, Rua Larga, Coimbra, Portugal

e-mail: iakh@isc-ras.ru

This abstract presents the main results of research on the conformational preference of the molecule mefenamic acid. This compound is used as a drug and belongs to the group of NSAID (nonsteroidal anti-inflammatory drugs). The main idea of this research is to determine the conformation preference of mefenamic acid dissolved in deuterated DMSO using two-dimensional NMR and quantum chemistry calculation. The obtained information can help one to determine the way of nucleation mechanisms.




Figure 1. Structure of the mefenamic acid polymorphs molecules MEF II (left) and MEF I (right).

Two crystalline forms, called MEF I and MEF II, having different positions of the carboxyl group, are formed by different conformations of the mefenamic acid molecules (Fig.1)[1].

Recently we have shown that the molecular conformation, existing in the saturated solution, may define the polymorphic type, obtained by evaporation of this solution [2,3].

In this work we present the conformation equilibrium of mefenamic acid in solution and discuss it in the frame of polymorphic phenomena.

**Acknowledgements.** The study was carried out with financial support of Russian Foundation for Basic Research (project 18-29-06008 MK), Ministry of Science and Higher Education and of the Russian Federation (projects 01201260481 and 01200950825 ), and under grant of Council on grants of the President of the Russian Federation (projects project nos. 18-29-06008, 18-03-00255 and 17-03-00459). Prof. V.V. Klochkov (from Kazan Federal University) and Prof. M.G. Kiselev (from ISC RAS) are appreciated for the helpful discussions of the NMR spectroscopy results.

[1] E.H. Lee, S.R. Byrn, and R. Pinal, *J. Pharm. Sci.* **101**, 4529 (2012).

[2] I.A. Khodov, S.V. Efimov, M.Y. Nikiforov, V.V. Klochkov, and N. Georgi, *J. Pharm. Sci.*, 103 (2014).

[3] I.A. Khodov, S.V. Efimov, M.G. Kiselev, L.A.E. Batista De Carvalho, and V.V. Klochkov, *J. Mol. Struct.*, 1113 (2016).