

## INTERIOR RADII OF SYMMETRIC NOT LEANING DOMAINS

L. V. Kovalyov

Many extremal problems for classes of analytic functions are reduced to problems on not leaning domains (see, e. g., [1], pp. 552–554; [2]; survey of new results in this direction can be found in [3]). Let  $D$  be a domain of the complex sphere  $\overline{\mathbb{C}}$ . We denote by  $g_D(z, z_0)$  Green's function of the domain  $D$  with the pole  $z_0$ ; the interior radius of  $D$  with respect to  $z_0$  is

$$r(D, z_0) \stackrel{\text{def}}{=} \begin{cases} \exp \left[ \lim_{z \rightarrow z_0} (g_D(z, z_0) + \log |z - z_0|) \right], & z_0 \neq \infty; \\ \exp \left[ \lim_{z \rightarrow z_0} (g_D(z, z_0) - \log |z|) \right], & z_0 = \infty. \end{cases}$$

Set  $D^* \stackrel{\text{def}}{=} \{z : 1/\bar{z} \in D\}$ .

The article is devoted to solving the following problem, posed in survey [4]. Let  $B_k$ ,  $k = 0, \dots, n$ , be pairwise not leaning domains in  $\overline{\mathbb{C}}$ ,  $a_k \in B_k$ . Find the least upper bound of the product  $\prod_{k=0}^n r(B_k, a_k)$  under the conditions  $a_0 = 0$ ,  $|a_k| = 1$ ,  $B_k = B_k^*$ ,  $k = 1, \dots, n$ . This statement strengthens Bakhtina's problem (see [5]), where the simple connection of domains  $B_k$  was supposed and the condition  $B_0 \subset \{z : |z| < 1\}$  was fulfilled; in these assumptions, in [5] the qualitative characteristic of the extremal configuration in terms of quadratic differentials was obtained (see [6], p. 48).

**Theorem.** *Let  $B_0, \dots, B_n$  ( $n > 2$ ) be not leaning domains in  $\overline{\mathbb{C}}$ ;  $a_k \in B_k$ ,  $k = 0, \dots, n$ ;  $a_0 = 0$ ,  $|a_k| = 1$ ,  $k = 1, \dots, n$ ;  $B_k = B_k^*$ ,  $k = 1, \dots, n$ . Then*

$$\prod_{k=0}^n r(B_k, a_k) \leq \frac{2^{2n+1/n}}{(n^2 - 2)^{n/2+1/n}} \left( \frac{n - \sqrt{2}}{n + \sqrt{2}} \right)^{\sqrt{2}}. \quad (1)$$

*If, in addition, the domains  $B_k$  possess classical Green's functions, then the equality in (1) is attained if and only if the points  $a_k$  and domains  $B_k$  are poles and circular domains, respectively, of the quadratic differential*

$$Q(z)dz^2 = -\frac{(\alpha z)^{2n} + (2n^2 - 2)(\alpha z)^n + 1}{z^2((\alpha z)^n - 1)^2} dz^2, \quad |\alpha| = 1.$$

Sketch of the proof. Let  $a_k = \exp(i\theta_k)$ ,  $0 = \theta_1 < \dots < \theta_n < \theta_{n+1} = 2\pi$ ,  $\varphi_k = \theta_{k+1} - \theta_k$ ,  $k = 1, \dots, n$ . Consider the two cases.

1. Suppose that  $\varphi_k \leq \pi\sqrt{2}$ ,  $k = 1, \dots, n$ . Let  $D_i = \{z : |z| < 1, \theta_i < \arg z < \theta_{i+1}\}$ . Let us apply a separating transformation (see [7]) of each domain  $B_k$  with respect to an appropriate family of functions which conformally map the domains  $D_i, D_i^*$ ,  $i = 1, \dots, n$ , onto a halfplane. By the same token the determination of the least upper bound of the left side of (1) is reduced to the estimation

---

Supported by Russian Foundation for Basic Research (grant 99-01-00443) and ISSEP (grant s97-336).

©2000 by Allerton Press, Inc.

Authorization to photocopy individual items for internal or personal use, or the internal or personal use of specific clients, is granted by Allerton Press, Inc. for libraries and other users registered with the Copyright Clearance Center (CCC) Transactional Reporting Service, provided that the base fee of \$ 50.00 per copy is paid directly to CCC, 222 Rosewood Drive, Danvers, MA 01923.