

# *Quantum Logics of Idempotents of Unital Rings*

**Airat Bikchentaev, Mirko Navara & Rinat Yakushev**

**International Journal of Theoretical Physics**

ISSN 0020-7748

Volume 54

Number 6

Int J Theor Phys (2015) 54:1987-2000

DOI 10.1007/s10773-014-2405-1

Volume 54 • Number 6 • June 2015

International  
Journal of  
Theoretical  
Physics

10773 • ISSN 0020-7748  
54(6) 1739–2118 (2015)

 Springer

 Springer

**Your article is protected by copyright and all rights are held exclusively by Springer Science +Business Media New York. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at [link.springer.com](http://link.springer.com)".**

# Quantum Logics of Idempotents of Unital Rings

Airat Bikchentaev · Mirko Navara · Rinat Yakushev

Received: 25 April 2014 / Accepted: 29 October 2014 / Published online: 18 November 2014  
© Springer Science+Business Media New York 2014

**Abstract** We introduce some new examples of quantum logics of idempotents in a ring. We continue the study of *symmetric logics*, i.e., collections of subsets generalizing Boolean algebras and closed under the symmetric difference.

**Keywords** Orthomodular poset · Quantum logic · State · Symmetric difference · Boolean algebra · Set representation ·  $C^*$ -algebra · Von Neumann algebra · Positive functional · Trace · Idempotent · Projection · Additive mapping

## 1 Motivation

Orthomodular posets and, in particular, orthomodular lattices appear as algebraic structures of events in quantum mechanics, cf. [14, 17, 31, 32]. The natural requirement that the event system must allow “sufficiently many” states leads (in its stronger form) to orthomodular posets which can be represented as collections of subsets of a set generalizing  $\sigma$ -algebras [14]. In such collections, the set-theoretical symmetric difference can be introduced as an additional operation [29] which cannot be derived from the lattice-theoretical

---

Dedicated to memory of Professor Daniar Mushtari

A. Bikchentaev · R. Yakushev  
N.I. Lobachevskii Institute of Mathematics and Mechanics, Kazan Federal University,  
Kremlevskaya 18, 420008, Kazan, Russian Federation  
e-mail: Airat.Bikchentaev@kpfu.ru

R. Yakushev  
e-mail: sultanich@rambler.ru

M. Navara (✉)  
Center for Machine Perception, Department of Cybernetics, Faculty of Electrical Engineering,  
Czech Technical University in Prague, Technická 2, 166 27 Prague 6, Czech Republic  
e-mail: navara@cmp.felk.cvut.cz  
URL: <http://cmp.felk.cvut.cz/~navara>

operations and orthocomplementation [21]. Thus we arrive at the notion of a symmetric logic.

During the study of symmetric logics, many questions remained open (cf. [5, 6]). In [7] we answered some of them. Here we present a generalization of [7, Theorem 4.3] with a shorter and direct proof.

## 2 Basic Notions

### 2.1 Quantum Logics of Idempotents of Unital Rings

**Definition 2.1** Let  $(L, \leq, 0, 1, \perp)$  be a poset with 0 and 1 as the smallest and greatest element, respectively, and a unary operation  $\perp : L \rightarrow L$  (the *orthocomplementation*) such that

- (i)  $p \leq q \Rightarrow q^\perp \leq p^\perp, \quad p, q \in L;$
- (ii)  $(p^\perp)^\perp = p, \quad p \in L;$
- (iii)  $p \vee p^\perp = 1, \quad p \in L;$
- (iv)  $p \leq q^\perp \Rightarrow p \vee q$  exists in  $L, \quad p, q \in L;$
- (v)  $p \leq q \Rightarrow q = p \vee (p^\perp \wedge q), \quad p, q \in L.$

Then  $L$  will be called a *quantum logic* or also an orthomodular poset. If  $L$  is also a lattice, then  $L$  is called an *orthomodular lattice*.

Let  $\mathcal{R}$  be a ring with unit  $e, x^\perp := e - x$  for  $\mathcal{R}$ . Then  $(x^\perp)^\perp = x$ . The set  $\mathcal{R}^{\text{id}} := \{x \in \mathcal{R} : x = x^2\}$ , equipped with the partial order  $p \leq q \Leftrightarrow pq = qp = p$  and orthocomplementation  $p \mapsto p^\perp$ , is a quantum logic. The logics  $\mathcal{R}^{\text{id}}$  are the main topic of this paper. They were investigated e.g. in [12, 13, 16, 18, 19, 25, 26].

**Definition 2.2** Let  $(L, \leq, 0, 1, \perp)$  be a quantum logic. A subset  $S$  of  $L$  is said to be a sublogic of  $L$  if the following conditions are satisfied:

- (i)  $0 \in S;$
- (ii) if  $p \in S$  then  $p^\perp \in S;$
- (iii) if  $p, q \in S$  and  $p \leq q^\perp$ , then  $p \vee q \in S.$

Let  $\mathcal{R}$  be an associative unital  $*$ -ring. Then the set  $\mathcal{R}^{\text{pr}} := \{x \in \mathcal{R} : x = x^* = x^2\}$  of all projections of  $\mathcal{R}$  is a sublogic of the logic  $\mathcal{R}^{\text{id}}$ . Let  $\langle \mathcal{R}, \|\cdot\| \rangle$  be a unital Banach  $*$ -algebra,  $\mathcal{R}_1 := \{x \in \mathcal{R} : \|x\| \leq 1\}$ . A linear functional  $\varphi$  on  $\mathcal{R}$  is called *positive* if  $\varphi(x^*x) \geq 0$  for every  $x \in \mathcal{R}$ . Every positive linear functional  $\varphi$  on  $\mathcal{R}$  is continuous and  $\|\varphi\| = \varphi(e)$  [34, Chap. I, Lemma 9.9]. A positive linear functional of norm one is called a *state* [34, Chap. I, Definition 9.4].

Let  $\mathcal{H}$  be a Hilbert space over  $\mathbb{C}$ , and  $\mathcal{B}(\mathcal{H})$  be the  $*$ -algebra of all bounded linear operators on  $\mathcal{H}$ . The *strong (operator) topology* on  $\mathcal{B}(\mathcal{H})$  is the locally convex topology determined by the seminorms  $x \in \mathcal{B}(\mathcal{H}) \mapsto \|x\xi\|_{\mathcal{H}}, \xi \in \mathcal{H}$ .

By the *commutant* of a set  $\mathcal{X} \subset \mathcal{B}(\mathcal{H})$  we mean the set

$$\mathcal{X}' = \{y \in \mathcal{B}(\mathcal{H}) : xy = yx, x^*y = yx^* \quad (x \in \mathcal{X})\}.$$

A  $*$ -subalgebra  $\mathcal{R}$  of the algebra  $\mathcal{B}(\mathcal{H})$  is called a *von Neumann algebra* acting in the Hilbert space  $\mathcal{H}$  if  $\mathcal{R} = \mathcal{R}''$ . A complex Banach  $*$ -algebra  $\mathcal{R}$  is called a *C\*-algebra* if

$\|x^*x\| = \|x\|^2$  for all  $x \in \mathcal{R}$ . Many  $C^*$ -algebras are generated as rings by their projections [1–4]. More precisely, every element in such a  $C^*$ -algebra  $\mathcal{R}$  can be represented as a finite sum of finite products of projections from  $\mathcal{R}$ .

For  $C^*$ -algebra  $\mathcal{R}$  let  $\mathcal{R}^+$  denote its positive part. A linear functional  $\varphi : \mathcal{R} \rightarrow \mathbb{C}$  is called a *trace* if  $\varphi(z^*z) = \varphi(zz^*)$  for all  $z \in \mathcal{R}$ . A positive linear functional  $\varphi$  on a von Neumann algebra  $\mathcal{R}$  is *normal* if  $x_i \nearrow x \implies \varphi(x) = \sup \varphi(x_i)$  ( $x_i, x \in \mathcal{R}^+$ ).

### 2.2 Concrete Logics

Let  $\Omega$  be a non-empty set. By  $2^\Omega$  we denote the set of all subsets of  $\Omega$ . For  $n \in \mathbb{N}$ , we define  $\Omega_n = \{1, 2, \dots, n\}$ .

Let us recall [14] that a collection  $\mathcal{E} \subseteq 2^\Omega$  of subsets of  $\Omega$  is called a *concrete (quantum) logic* if the following conditions hold true:

- (C1)  $\Omega \in \mathcal{E}$ ,
- (C2)  $A \in \mathcal{E} \implies A^c := \Omega \setminus A \in \mathcal{E}$ ,
- (C3)  $A, B \in \mathcal{E}, A \cap B = \emptyset \implies A \cup B \in \mathcal{E}$ .

A concrete logic  $\mathcal{E}$  is called a  $\sigma$ -class [14] if it satisfies the following strengthening of (C3):

- (C3')  $\{A_n \mid n \in \mathbb{N}\} \subseteq \mathcal{E}, A_m \cap A_n = \emptyset$  whenever  $m \neq n \implies \bigcup_{n \in \mathbb{N}} A_n \in \mathcal{E}$ .

A family  $\mathcal{E} \subseteq 2^\Omega$  is a concrete logic if and only if it satisfies (C1) and the following condition:

- (C4)  $A, B \in \mathcal{E}, A \subseteq B \implies B \setminus A \in \mathcal{E}$ .

*Remark 2.3* Every concrete logic can be represented as the logic of idempotents in some ring. Let  $\Omega$  be a non-empty set, and let  $\mathcal{E} \subseteq 2^\Omega$  be a concrete logic. If  $\mathbb{R}^\Omega$  is the ring of all real functions on  $\Omega$ , then the set of all characteristic functions  $\chi_A, A \in \mathcal{E}$ , is a logic of idempotents of  $\mathbb{R}^\Omega$ . This logic is isomorphic to  $\mathcal{E}$ .

### 2.3 Symmetric Logics

The set  $2^\Omega$  is a group with respect to the symmetric difference operation:  $A \Delta B := (A \setminus B) \cup (B \setminus A)$ . Notice that

$$\begin{aligned} A^c \Delta B &= (A \Delta B)^c, \\ A^c \Delta B^c &= A \Delta B. \end{aligned}$$

A *symmetric logic* [28, Definition 3.2] is a concrete quantum logic  $\mathcal{E}$  satisfying:

- (S)  $A, B \in \mathcal{E} \implies A \Delta B \in \mathcal{E}$ .

A family  $\mathcal{E} \subseteq 2^\Omega$  is a symmetric logic if and only if it satisfies (C1) and (S) [5, Proposition 1]. Symmetric logics were investigated e.g. in [5, 6, 10, 11, 21, 22, 28, 29].

*Example 2.4* Let  $n \in \mathbb{N}$  and  $\Omega_{2n} = \{1, 2, \dots, 2n\}$ . Then the family

$$\mathcal{E}_{2n}^{\text{even}} = \{A \subseteq \Omega_{2n} \mid \text{card } A \text{ is even}\}$$

is a symmetric logic on  $\Omega_{2n}$ .

*Example 2.5* Let  $\mathcal{E} \subset 2^\Omega$  be a concrete quantum logic and  $T \in \mathcal{E}$ ,  $T \neq \emptyset$ . Then the family  $\mathcal{E}_T = \{A \in \mathcal{E} \mid A \subseteq T\}$  is a concrete quantum logic with the greatest element  $T$ . Moreover, if  $\mathcal{E}$  is a symmetric logic, then  $\mathcal{E}_T$  is also a symmetric logic.

In the latter example, it was necessary to assume that  $T \in \mathcal{E}$ . This condition can be omitted in symmetric logics.

*Example 2.6* Let  $\mathcal{E} \subseteq 2^\Omega$  be a symmetric logic and  $T \subseteq \Omega$ ,  $T \neq \emptyset$ . Then the family

$$\mathcal{E}|_T = \{A \cap T \mid A \in \mathcal{E}\} \subseteq 2^T$$

is a symmetric logic with the greatest element  $T$ .

## 2.4 States

We say that a mapping  $m : \mathcal{E} \rightarrow [0, 1]$  is a *state* (or a finitely additive *probability measure*) on a concrete logic  $\mathcal{E}$  if  $m(\Omega) = 1$  and  $m(A \cup B) = m(A) + m(B)$  whenever  $A, B \in \mathcal{E}$ ,  $A \cap B = \emptyset$ . Let us denote by  $P(\mathcal{E})$  the set of all states on a concrete logic  $\mathcal{E}$ . Recall that a state  $m \in P(\mathcal{E})$  is called *subadditive* [31, p. 829] if for each  $A, B \in \mathcal{E}$  there exists a set  $C \in \mathcal{E}$  such that  $C \supseteq A \cup B$  and  $m(C) \leq m(A) + m(B)$ .

If  $\mathcal{E}$  is a Boolean algebra then any state  $m \in P(\mathcal{E})$  is subadditive. There exists a concrete quantum logic which is not a Boolean algebra and all of its states are subadditive. This result was established in [30] with substantial help of the techniques developed in [23] and [27] (see also [31, p. 831]).

From now on, we suppose that  $\mathcal{E}$  is a symmetric logic. A state  $m \in P(\mathcal{E})$  is called  $\Delta$ -subadditive [10] if

$$m(A \Delta B) \leq m(A) + m(B) \text{ for any pair } A, B \in \mathcal{E}.$$

The set of all  $\Delta$ -subadditive states is convex. Every subadditive state  $m \in P(\mathcal{E})$  is  $\Delta$ -subadditive (hint:  $C \supseteq A \cup B \supseteq A \Delta B$ ), but the reverse implication does not hold in general. In [6], the following situations were demonstrated:

- 1) a  $\Delta$ -subadditive state which is not subadditive;
- 2) a two-valued state which is not  $\Delta$ -subadditive.

## 3 Additive Mappings and Quantum Logics

### 3.1 New Quantum Logics of Idempotents in a Ring

**Theorem 3.1** *Let  $\mathcal{R}$  be a ring with unit  $e$ ;  $x, y \in \mathcal{R}$ , and  $\varphi : \mathcal{R} \rightarrow \mathbb{C}$  be an additive mapping with  $\varphi(e) = 1$ . Then the sets*

$$\mathcal{R}_{\varphi,1}^{x,y} := \{p \in \mathcal{R}^{\text{id}} : \varphi(px + yp) = \varphi(p)\varphi(x + y)\}$$

and

$$\mathcal{R}_{\varphi,2}^{x,y} := \{p \in \mathcal{R}^{\text{id}} : \varphi(xpy) = \varphi(p)\varphi(xy)\}$$

are quantum logics with the greatest element  $e$ , the partial order inherited from  $\mathcal{R}^{\text{id}}$  and the orthocomplementation  $p \mapsto p^\perp = e - p$ .

Moreover, if  $\langle \mathcal{R}, t \rangle$  is a topological ring and  $\varphi$  is  $t$ -continuous, then the sets  $\mathcal{R}_{\varphi,1}^{x,y}$  and  $\mathcal{R}_{\varphi,2}^{x,y}$  are  $t$ -closed.

*Proof* It is clear that  $0, e \in \mathcal{R}_{\varphi,k}^{x,y}$  for  $k \in \{1, 2\}$ . We show that  $p \in \mathcal{R}_{\varphi,k}^{x,y} \Leftrightarrow p^\perp \in \mathcal{R}_{\varphi,k}^{x,y}$  for all  $p \in \mathcal{R}^{\text{id}}$  and  $k \in \{1, 2\}$ . Let  $p \in \mathcal{R}_{\varphi,1}^{x,y}$ . Since  $p^\perp x + y p^\perp = x + y - (px + yp)$ , we have  $\varphi(p^\perp x + y p^\perp) = \varphi(x + y) - \varphi(px + yp) = \varphi(x + y) - \varphi(p)\varphi(x + y) = \varphi(p^\perp)\varphi(x + y)$  and  $p^\perp \in \mathcal{R}_{\varphi,1}^{x,y}$ . Let now  $p \in \mathcal{R}_{\varphi,2}^{x,y}$ . Since  $x p^\perp y = xy - xpy$ , we have

$$\varphi(x p^\perp y) = \varphi(xy) - \varphi(xpy) = \varphi(xy) - \varphi(p)\varphi(xy) = \varphi(p^\perp)\varphi(xy)$$

and  $p^\perp \in \mathcal{R}_{\varphi,2}^{x,y}$ .

Let  $p, q \in \mathcal{R}_{\varphi,k}^{x,y}$  for  $k \in \{1, 2\}$ .

If  $p \leq q^\perp$ , then  $p \vee q = p + q \in \mathcal{R}^{\text{id}}$  and it is easy to check that  $p \vee q \in \mathcal{R}_{\varphi,k}^{x,y}$ .

If  $p \leq q$ , then  $q - p \in \mathcal{R}^{\text{id}}$ ,  $q - p \leq p^\perp$ , and  $q = (q - p) \vee p$ . It is easy to check that  $q - p \in \mathcal{R}_{\varphi,k}^{x,y}$ .

Finally, note that if  $\langle \mathcal{R}, t \rangle$  is a topological ring, then the quantum logic  $\mathcal{R}^{\text{id}}$ , being defined by equalities containing continuous operations, is  $t$ -closed.  $\square$

**Proposition 3.2** *Let  $x, y, u, v \in \mathcal{R}$  and  $p, q \in \mathcal{R}^{\text{id}}$ . Then the following holds:*

- 1)  $\mathcal{R}_{\varphi,1}^{0,0} = \mathcal{R}_{\varphi,1}^{e,0} = \mathcal{R}_{\varphi,1}^{0,e} = \mathcal{R}_{\varphi,1}^{e,e} = \mathcal{R}_{\varphi,2}^{x,0} = \mathcal{R}_{\varphi,2}^{0,y} = \mathcal{R}_{\varphi,2}^{e,e} = \mathcal{R}^{\text{id}}$ .
- 2)  $\lambda, \mu \in \mathbb{Z} \implies \mathcal{R}_{\varphi,1}^{\lambda e \pm x, \mu e \pm y} = \mathcal{R}_{\varphi,1}^{x,y}$  for the following choices of signs in two  $\pm$ :  $+, +$  and  $-, -$ .
- 3)  $\mathcal{R}_{\varphi,k}^{-x,-y} = \mathcal{R}_{\varphi,k}^{x,y}$  for  $k \in \{1, 2\}$ .
- 4)  $\mathcal{R}_{\varphi,1}^{x,y} \cap \mathcal{R}_{\varphi,1}^{u,v} \subset \mathcal{R}_{\varphi,1}^{x+u,y+v}$ .
- 5)  $\mathcal{R}_{\varphi,1}^{x,0} = \mathcal{R}_{\varphi,2}^{e,x}$ .
- 6)  $\mathcal{R}_{\varphi,1}^{0,y} = \mathcal{R}_{\varphi,2}^{y,e}$ .
- 7)  $p \in \mathcal{R}_{\varphi,1}^{q,0} \Leftrightarrow q \in \mathcal{R}_{\varphi,1}^{0,p}$ .
- 8)  $p \in \mathcal{R}_{\varphi,1}^{q,q} \Leftrightarrow q \in \mathcal{R}_{\varphi,1}^{p,p}$ .
- 9)  $p \in \mathcal{R}_{\varphi,1}^{p,p} \Leftrightarrow p \in \mathcal{R}_{\varphi,2}^{p,p} \Leftrightarrow \varphi(p) \in \{0, 1\}$ .

*Proof* 1) Easy verification.

2) We have

$$p(\lambda e \pm x) + (\mu e \pm y)p = (\lambda + \mu)p \pm (px + yp), \tag{1}$$

i.e.  $\mp(px + yp) = (\lambda + \mu)p - (p(\lambda e \pm x) + (\mu e \pm y)p)$ . The inclusion “ $\subset$ ”:

$$\begin{aligned} \mp\varphi(px + yp) &= (\lambda + \mu)\varphi(p) - \varphi(p(\lambda e \pm x) + (\mu e \pm y)p) \\ &= (\lambda + \mu)\varphi(p) - \varphi(p)\varphi((\lambda + \mu)e \pm (x + y)) \\ &= (\lambda + \mu)\varphi(p) - \varphi(p)(\lambda + \mu \pm \varphi(x + y)) = \mp\varphi(p)\varphi(x + y). \end{aligned}$$

The inclusion “ $\supset$ ”: we have via (1)

$$\begin{aligned} \varphi(p(\lambda e \pm x) + (\mu e \pm y)p) &= \varphi((\lambda + \mu)p \pm (px + yp)) = (\lambda + \mu)\varphi(p) \pm \varphi(px + yp) \\ &= (\lambda + \mu)\varphi(p) \pm \varphi(p)\varphi(x + y) \\ &= \varphi(p)\varphi((\lambda e \pm x) + (\mu e \pm y)). \end{aligned}$$

- 3) For  $k = 1$ , it follows by 2) with  $\lambda = \mu = 0$ . For  $k = 2$  we have  $\varphi((-x)p(-y)) = \varphi(p)\varphi((-x)(-y)) \Leftrightarrow \varphi(xpy) = \varphi(p)\varphi(xy)$ .
- 5) We have  $\varphi(px) = \varphi(p)\varphi(x) \Leftrightarrow \varphi(epx) = \varphi(p)\varphi(ex)$ .
- 6) We have  $\varphi(yq) = \varphi(p)\varphi(y) \Leftrightarrow \varphi(yqe) = \varphi(p)\varphi(ye)$ .
- 7) We have  $\varphi(pq + 0p) = \varphi(p)\varphi(q) \Leftrightarrow \varphi(q0 + pq) = \varphi(q)\varphi(p)$ .
- 8) We have  $\varphi(pq + qp) = \varphi(p)\varphi(2q) \Leftrightarrow \varphi(qp + pq) = \varphi(q)\varphi(2p)$ .
- 9) We have  $2\varphi(p) = \varphi(pp + pp) = \varphi(p)\varphi(p + p) \Leftrightarrow \varphi(p) = (\varphi(p))^2 \Leftrightarrow \varphi(p) \in \{0, 1\}$  and  $\varphi(ppp) = \varphi(p)\varphi(pp) \Leftrightarrow \varphi(p) = (\varphi(p))^2 \Leftrightarrow \varphi(p) \in \{0, 1\}$ .

□

**Remark 3.3** We obtain  $\mathcal{R}_{\varphi,1}^{u,v} \cap \mathcal{R}_{\varphi,1}^{u+x,v+y} \subset \mathcal{R}_{\varphi,1}^{x,y}$  by 3) and 4) of Proposition 3.2. If  $\mathcal{R}$  is a unital algebra, then  $\mathcal{R}_{\varphi,1}^{\lambda e \pm x, \mu e \pm y} = \mathcal{R}_{\varphi,1}^{x,y}$  for all  $\lambda, \mu \in \mathbb{C}$  and for the following choices of signs in two  $\pm$ : +, + and -, -.

**Proposition 3.4** Let  $t \in \mathcal{R}$  be invertible,  $\psi(z) := \varphi(tzt^{-1})$  for all  $z \in \mathcal{R}$  and let  $p \in \mathcal{R}^{\text{id}}$ . Then  $p \in \mathcal{R}_{\psi,k}^{x,y} \Leftrightarrow tpt^{-1} \in \mathcal{R}_{\varphi,k}^{txt^{-1},tyt^{-1}}$  for all  $x, y \in \mathcal{R}$  and  $k \in \{1, 2\}$ .

*Proof* The implication “ $\Rightarrow$ ”: If  $k = 1$ , then

$$\begin{aligned} \varphi\left(tpt^{-1}txt^{-1} + tyt^{-1}tpt^{-1}\right) &= \varphi\left(t(px + yp)t^{-1}\right) = \psi(px + yp) = \psi(p)\psi(x + y) \\ &= \varphi\left(tpt^{-1}\right)\varphi\left(txt^{-1} + tyt^{-1}\right). \end{aligned}$$

If  $k = 2$ , then

$$\begin{aligned} \varphi\left(txt^{-1}tpt^{-1}tyt^{-1}\right) &= \varphi\left(txpyt^{-1}\right) = \psi(xpy) = \psi(p)\psi(xy) \\ &= \varphi\left(tpt^{-1}\right)\varphi\left(txt^{-1}tyt^{-1}\right). \end{aligned}$$

□

**Proposition 3.5** Let  $x, y \in \mathcal{R}$  and  $p \in \mathcal{R}^{\text{id}}$ . If  $py = yp$ , then

- 1)  $p \in \mathcal{R}_{\varphi,1}^{x,y} \Leftrightarrow p \in \mathcal{R}_{\varphi,1}^{x+y,0}$ ;
- 2)  $p \in \mathcal{R}_{\varphi,2}^{x,y} \Leftrightarrow p \in \mathcal{R}_{\varphi,1}^{0,xy}$ .

In particular, if  $y$  is a central element of  $\mathcal{R}$ , then  $\mathcal{R}_{\varphi,1}^{x,y} = \mathcal{R}_{\varphi,1}^{x+y,0}$  and  $\mathcal{R}_{\varphi,2}^{x,y} = \mathcal{R}_{\varphi,1}^{0,xy}$ .

### 3.2 Quantum Logics of Idempotents of Unital Banach \*-algebras

**Proposition 3.6** Let  $\langle \mathcal{R}, \|\cdot\| \rangle$  be a unital Banach \*-algebra,  $x, y \in \mathcal{R}$  and  $\varphi$  be a state on  $\mathcal{R}$ ,  $k \in \{1, 2\}$ .

- 1) The quantum logic  $\mathcal{R}_{\varphi,k}^{x,y}$  is  $\|\cdot\|$ -closed.
- 2)  $p \in \mathcal{R}_{\varphi,k}^{x,y} \Leftrightarrow p^* \in \mathcal{R}_{\varphi,k}^{y^*,x^*}$  for all  $p \in \mathcal{R}^{\text{id}}$ .

*Proof* 1) The quantum logic  $\mathcal{R}^{\text{id}}$  is  $\|\cdot\|$ -closed. Every positive linear functional on any unital Banach \*-algebra automatically is continuous [34, Chap. I, Lemma 9.9]. Hence the quantum logic  $\mathcal{R}_{\varphi,k}^{x,y}$  is  $\|\cdot\|$ -closed via Theorem 3.1.



2) Recall that  $(x^*)^* = x$  and  $(xy)^* = y^*x^*$ . We have  $\varphi(z^*) = \overline{\varphi(z)}$  for all  $z \in \mathcal{R}$  [34, Chap. I, §9, formula (3)]. If  $p \in \mathcal{R}_{\varphi,1}^{x,y}$ , then

$$\varphi(p^*y^* + x^*p^*) = \varphi((px + yp)^*) = \overline{\varphi(px + yp)} = \overline{\varphi(p) \cdot \varphi(x + y)} = \varphi(p^*)\varphi(x^* + y^*)$$

and  $p^* \in \mathcal{R}_{\varphi,1}^{y^*,x^*}$ . If  $p \in \mathcal{R}_{\varphi,2}^{x,y}$ , then

$$\varphi(y^*p^*x^*) = \varphi((xpy)^*) = \overline{\varphi(xpy)} = \overline{\varphi(p) \cdot \varphi(xy)} = \varphi(p^*)\varphi(y^*x^*)$$

and  $p^* \in \mathcal{R}_{\varphi,2}^{y^*,x^*}$ .

In particular, for  $y = x^*$  we have  $p \in \mathcal{R}_{\varphi,k}^{x,x^*} \Leftrightarrow p^* \in \mathcal{R}_{\varphi,k}^{x^*,x}$  for all  $p \in \mathcal{R}^{\text{id}}$  and  $k \in \{1, 2\}$ . □

**Theorem 3.7** *Let  $\mathcal{R}$  be an unital  $C^*$ -algebra,  $p \in \mathcal{R}^{\text{id}}$  and  $x \in \mathcal{R}$ . Then the following conditions are equivalent:*

- (i)  $xp = px$ ;
- (ii)  $p \in \mathcal{R}_{\varphi,1}^{x,e-x}$  for all states  $\varphi$  on  $\mathcal{R}$ .

*Proof* (ii) $\Rightarrow$ (i). We have  $\|\varphi\| = \varphi(e) = 1$  and  $\varphi(xp) = \varphi(px)$  for all states  $\varphi$  on  $\mathcal{R}$ . By Hahn-Banach separation theorem, the set  $\mathcal{R}^*$  of all continuous linear functionals on  $\mathcal{R}$  is separating for  $\mathcal{R}$ . If  $f \in \mathcal{R}^*$ , we define  $f^* \in \mathcal{R}^*$  by setting  $f^*(a) = \overline{f(a^*)}$  for all  $a \in \mathcal{R}$ . We say a functional  $f \in \mathcal{R}^*$  is *self-adjoint* if  $f = f^*$ . For any bounded linear functional  $f$  on  $\mathcal{R}$ , there are unique self-adjoint bounded linear functionals  $f_1$  and  $f_2$  on  $\mathcal{R}$  such that  $f = f_1 + if_2$  (take  $f_1 = (f + f^*)/2$  and  $f_2 = (f - f^*)/(2i)$ ). Let  $\tau$  be a self-adjoint bounded linear functional on  $C^*$ -algebra  $\mathcal{R}$ . Then by Jordan Decomposition Theorem [24, Theorem 3.3.10] there exist positive linear functionals  $\tau_+, \tau_-$  on  $\mathcal{R}$  such that  $\tau = \tau_+ - \tau_-$  and  $\|\tau\| = \|\tau_+\| + \|\tau_-\|$ . Thus every  $f \in \mathcal{R}^*$  is a linear combination of four positive ones. Hence, the set of all states on  $\mathcal{R}$  is separating for  $\mathcal{R}$  and  $xp = px$ . □

**Proposition 3.8** *Let a state  $\varphi$  on a von Neumann algebra  $\mathcal{R}$  be normal,  $x, y \in \mathcal{R}$  and  $k \in \{1, 2\}$ . Then the quantum logic  $\mathcal{R}_{\varphi,k}^{x,y} \cap \mathcal{R}^{\text{pr}}$  is so-closed.*

*Proof* Since  $\mathcal{B}(\mathcal{H})^{\text{pr}}$  is closed in the strong operator topology (i.e., so-closed) [15, Exercise 5.7.8] and  $\mathcal{R}$  is so-closed, the set  $\mathcal{R}^{\text{pr}} = \mathcal{B}(\mathcal{H})^{\text{pr}} \cap \mathcal{R}$  is so-closed. The multiplication operation  $(u, v) \mapsto uv$  is so-continuous as a mapping  $\mathcal{B}(\mathcal{H})_1 \times \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$  [8, Chap. II, Proposition 2.4.1]. Finally, recall that every normal state  $\varphi$  on a von Neumann algebra  $\mathcal{R}$  is so-continuous on  $\mathcal{R}_1$  [34, Chap. II, Theorem 2.6]. □

**Proposition 3.9** *If a state  $\varphi$  on a von Neumann algebra  $\mathcal{R}$  is singular, then for every nonzero  $p \in \mathcal{R}^{\text{pr}}$  there exists a nonzero  $q \in \mathcal{R}^{\text{pr}}$  such that  $q \leq p$  and  $q \in \mathcal{R}_{\varphi,1}^{p,0} \cap \mathcal{R}_{\varphi,1}^{0,p} \cap \mathcal{R}_{\varphi,1}^{p,p} \cap \mathcal{R}_{\varphi,2}^{p,p}$ .*

*Proof* For singular state  $\varphi$  for every nonzero  $p \in \mathcal{R}^{\text{pr}}$  there exists a nonzero  $q \in \mathcal{R}^{\text{pr}}$  such that  $q \leq p$  and  $\varphi(q) = 0$  [34, Chap. III, Theorem 3.8]. We have  $pq = qp = \frac{1}{2}(pq + qp) = pqp = q$  and

$$\varphi(pq) = \varphi(qp) = \varphi(pq + qp) = \varphi(pqp) = \varphi(q) = 0 = \varphi(q)\varphi(p). \quad \square$$

3.3 Quantum Logics and Tracial States on Unital  $C^*$ -algebras

**Proposition 3.10** *Let  $\varphi$  be a tracial state on unital  $C^*$ -algebra  $\mathcal{R}$  and  $k \in \{1, 2\}$ . Then the following holds:*

- 1)  $\mathcal{R}_{\varphi,2}^{x,y} = \mathcal{R}_{\varphi,1}^{yx,0}$  for all  $x, y \in \mathcal{R}$ .
- 2)  $\mathcal{R}_{\varphi,1}^{x,y} = \mathcal{R}_{\varphi,2}^{x+y,e}$  for all  $x, y \in \mathcal{R}$ .
- 3)  $\mathcal{R}_{\varphi,2}^{x,y} = \mathcal{R}^{\text{id}}$  for all  $x, y \in \mathcal{R}$  with  $yx \in \{0, e\}$ .
- 4)  $\mathcal{R}_{\varphi,1}^{\lambda e \pm x, \mu e \mp x} = \mathcal{R}^{\text{id}}$  for all  $x \in \mathcal{R}$  and  $\lambda, \mu \in \mathbb{C}$  (the signs in the formula must be opposite to each other).
- 5)  $\mathcal{R}_{\varphi,1}^{x,x} = \mathcal{R}_{\varphi,2}^{x,x}$  for all  $x \in \mathcal{R}^{\text{id}}$ .
- 6)  $\mathcal{R}_{\varphi,k}^{x,x^\perp} = \mathcal{R}^{\text{id}}$  for all  $x \in \mathcal{R}^{\text{id}}$ .
- 7)  $p \in \mathcal{R}_{\varphi,k}^{x,y} \Leftrightarrow tpt^{-1} \in \mathcal{R}_{\varphi,k}^{txt^{-1}, tyt^{-1}}$  for all  $p \in \mathcal{R}^{\text{id}}$ ,  $x, y \in \mathcal{R}$  and an invertible  $t \in \mathcal{R}$ .

*Proof* 1) The inclusion “ $\subset$ ”: we have  $\varphi(pyx) = \varphi(xpy) = \varphi(p)\varphi(xy) = \varphi(p)\varphi(yx)$ .  
 The inclusion “ $\supset$ ”: we have  $\varphi(xpy) = \varphi(pyx) = \varphi(p)\varphi(yx) = \varphi(p)\varphi(xy)$ .  
 2) The inclusion “ $\subset$ ”: we have  $\varphi(p)\varphi(x+y) = \varphi(px+yp) = \varphi(px) + \varphi(y) = \varphi((x+y)p) = \varphi((x+y)pe)$ . The inclusion “ $\supset$ ”: we have  $\varphi(px+yp) = \varphi(px) + \varphi(y) = \varphi((x+y)p) = \varphi((x+y)pe)$ .  
 3) Let  $p \in \mathcal{R}^{\text{id}}$ . If  $yx = 0$ , then  $0 = \varphi(pyx) = \varphi(xpy) = \varphi(p)\varphi(xy) = \varphi(p)\varphi(yx)$ . If  $yx = e$ , then  $\varphi(xpy) = \varphi(pyx) = \varphi(p) = \varphi(p)\varphi(yx) = \varphi(p)\varphi(xy)$ .  
 4) We have

$$\begin{aligned} \varphi(p(\lambda e \pm x) + (\mu e \mp x)p) &= \varphi((\lambda + \mu)p \pm (px - xp)) \\ &= (\lambda + \mu)\varphi(p) \pm (\varphi(px) - \varphi(xp)) \\ &= (\lambda + \mu)\varphi(p) = \varphi(p)\varphi((\lambda e \pm x) + (\mu e \mp x)) \end{aligned}$$

for all  $p \in \mathcal{R}^{\text{id}}$ .

5) The inclusion “ $\subset$ ”: we have  $\varphi(px+xp) = \varphi(px) + \varphi(xp) = 2\varphi(px) = \varphi(p)\varphi(2x) \Rightarrow \varphi(xpx) = \varphi(px^2) = \varphi(px) = \varphi(p)\varphi(x^2)$ .

The inclusion “ $\supset$ ”: we have  $\varphi(xpx) = \varphi(px^2) = \varphi(p)\varphi(x^2) = \varphi(p)\varphi(x) \Rightarrow \varphi(px+xp) = \varphi(px) + \varphi(xp) = 2\varphi(xpx) = 2\varphi(p)\varphi(x^2) = 2\varphi(p)\varphi(x) = \varphi(p)\varphi(x+x)$ .

6) Let  $p \in \mathcal{R}^{\text{id}}$ . If  $k = 1$ , then

$$\varphi(px + x^\perp p) = \varphi(px) + \varphi(x^\perp p) = \varphi(px + px^\perp) = \varphi(p) = \varphi(p)\varphi(x + x^\perp).$$

If  $k = 2$ , then  $\varphi(xpx^\perp) = \varphi(px^\perp x) = \varphi(0) = 0 = \varphi(p)\varphi(xx^\perp)$ .

7) We apply Proposition 3.4 with  $\psi = \varphi$ . □

*Example 3.11* Let  $\mathcal{R} = \mathbb{M}_2(\mathbb{C})$  and  $\varphi$  be the normalized trace on  $\mathcal{R}$ , i.e.  $\varphi\left(\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}\right) = \frac{1}{2}(\alpha + \delta)$ ,  $0 = \text{diag}(0, 0)$ ,  $e = \text{diag}(1, 1)$ . Put  $p(a, b, c) = \begin{pmatrix} a & b \\ c & 1-a \end{pmatrix}$  for  $a, b, c \in \mathbb{C}$  with  $a = a^2 + bc$ , then

$$\mathcal{R}^{\text{id}} = \{0, e, p(a, b, c) \text{ with } a, b, c \in \mathbb{C} \text{ and } a = a^2 + bc\}$$

is a quantum logic which is a lattice. For  $x = p(1, 0, 0)$  and  $y = p(1/2, 1/2, 1/2)$  we have

$$\mathcal{R}_{\varphi,1}^{x,y} = \{0, e, p(a, b, c), \text{ where } a, b, c \in \mathbb{C} \text{ with } a = a^2 + bc \text{ and } 2a + b + c = 1\},$$

$$\mathcal{R}_{\varphi,2}^{x,y} = \{0, e, p(a, b, c), \text{ where } a, b, c \in \mathbb{C} \text{ with } a = a^2 + bc \text{ and } 2a + 2b = 1\}.$$

Hence  $\mathcal{R}_{\varphi,1}^{x,y} \cap \mathcal{R}_{\varphi,2}^{x,y} = \left\{0, e, q = p\left(\frac{1}{2} - \frac{1}{2^{3/2}}, \frac{1}{2^{3/2}}, \frac{1}{2^{3/2}}\right), q^\perp\right\}$ . Also we have

$$p(0, 1, 0) \in \mathcal{R}_{\varphi,1}^{x,y} \setminus \mathcal{R}_{\varphi,2}^{x,y}, \quad p(1/4, 1/4, 3/4) \in \mathcal{R}_{\varphi,2}^{x,y} \setminus \mathcal{R}_{\varphi,1}^{x,y}.$$

### 4 Concrete Quantum Logics

#### 4.1 Asymmetric Logics: Definition and Examples

**Definition 4.1** A concrete logic  $\mathcal{E}$  is called an *asymmetric logic* if  $A \Delta B \in \mathcal{E}$  if and only if  $A \cap B \in \mathcal{E}$  for all  $A, B \in \mathcal{E}$ .

*Example 4.2* Let  $\Omega = \{z_n\}_{n=1}^\infty$  be a sequence of complex numbers such that  $\Omega \in \ell_1$ , i.e. the series  $\sum_{n=1}^\infty z_n$  converges absolutely. Let  $\Lambda \in \{\mathbb{Q}, \mathbb{R}\}$  and  $z = \sum_{n=1}^\infty z_n$ . Recall that every rearrangement of  $\{z_n\}_{n=1}^\infty$  preserves the absolute convergence and the sum  $z$ . Then

$$\mathcal{E}_{\Lambda,\Omega} = \{I \subset \Omega \mid \sum_{x \in I} x = \lambda z \text{ for some } \lambda \in \Lambda\}$$

is an asymmetric logic. (The sum of an empty sequence is considered zero, thus  $\emptyset \in \mathcal{E}_{\Lambda,\Omega}$ .) Moreover,  $\mathcal{E}_{\mathbb{R},\Omega}$  is a  $\sigma$ -class and  $\mathcal{E}_{\mathbb{Q},\Omega}$  is its sublogic.

*Example 4.3* Let  $\mathcal{A}$  be the Lebesgue  $\sigma$ -algebra on  $\Omega = [0, 1]$ ,  $\mu$  be the linear Lebesgue measure such that  $\mu(\Omega) = 1$ . Then  $\mathcal{E}_{\mathbb{Q},\mu} = \{A \in \mathcal{A} : \mu(A) \in \mathbb{Q}\}$  is an asymmetric logic.

Symmetric logics may be asymmetric, e.g., Boolean algebras, or may not be asymmetric, e.g.  $\mathcal{E}_4^{\text{even}}$ . The latter example is prototypical in the following sense:

**Proposition 4.4** *If  $\mathcal{E}$  is a symmetric logic of subsets of  $\Omega$  and  $\mathcal{E}$  is not an asymmetric logic, then there is a partition  $\{C_i\}_{i=1}^4$  of  $\Omega$  with the following property:*

*For  $I \subset \{1, 2, 3, 4\}$ , the union  $\bigcup_{i \in I} C_i$  belongs to  $\mathcal{E}$  if and only if  $\text{card } I$  is even.*

*Proof* If  $\mathcal{E}$  is not an asymmetric logic, then there are  $A, B \in \mathcal{E}$  such that  $A \Delta B \in \mathcal{E}$  and  $A \cap B \notin \mathcal{E}$ . It suffices to take  $C_1 = A \cap B^c, C_2 = A^c \cap B, C_3 = A \cap B, C_4 = A^c \cap B^c$ .  $\square$

**Proposition 4.5** *A symmetric logic is an asymmetric logic if and only if it is a Boolean algebra.*

Together with Proposition 4.4, we obtain:

**Corollary 4.6** *If a symmetric logic is not a Boolean algebra, it contains a sublogic isomorphic to  $\mathcal{E}_4^{\text{even}}$ .*

### 4.2 Concrete Logics Generated by the Independence Relation

Let  $\mathcal{A}$  be a Boolean algebra with the unit  $\Omega$ ,  $\varphi : \mathcal{A} \rightarrow \mathbb{C}$  be an additive mapping ( $\varphi(A \cup B) = \varphi(A) + \varphi(B)$  for all  $A, B \in \mathcal{A}$ ,  $A \cap B = \emptyset$ ) with  $\varphi(\Omega) = 1$ . Let  $A, B \in \mathcal{A}$ . We have  $\varphi(A) + \varphi(A^c) = \varphi(\Omega) = 1$  and  $\varphi(A^c) = 1 - \varphi(A)$ , hence  $\varphi(\emptyset) = 0$ . The following conditions are equivalent:

- (i)  $\varphi(A \cap B) = \varphi(A)\varphi(B)$ ;
- (ii)  $\varphi(A^c \cap B) = \varphi(A^c)\varphi(B)$ ;
- (iii)  $\varphi(A \cap B^c) = \varphi(A)\varphi(B^c)$ ;
- (iv)  $\varphi(A^c \cap B^c) = \varphi(A^c)\varphi(B^c)$ .

**Proposition 4.7** *The family*

$$\mathcal{A}_\varphi^A := \{B \in \mathcal{A} : \varphi(A \cap B) = \varphi(A)\varphi(B)\}$$

*is a concrete logic with the greatest element  $\Omega$ . We have  $\mathcal{A}_\varphi^A = \mathcal{A}_\varphi^{A^c}$ . Moreover, if  $\mathcal{A}$  is a  $\sigma$ -algebra and  $\varphi$  is  $\sigma$ -additive, then  $\mathcal{A}_\varphi^A$  is a  $\sigma$ -class.*

*Proof* It follows by distributivity of the intersection with respect to the union. □

Let  $\mathcal{A}$  be a Boolean algebra and  $\nu : \mathcal{A} \rightarrow \mathbb{R}$  be a measure ( $\nu(A \cup B) = \nu(A) + \nu(B)$  for all  $A, B \in \mathcal{A}$ ,  $A \cap B = \emptyset$ ). An event  $A \in \mathcal{A}$  is a  $\nu$ -atom if  $\nu(A) > 0$  and if for any event  $B \subset A$ , either  $\nu(B) = \nu(A)$  or  $\nu(B) = 0$ . A measure  $\nu$  is *nonatomic* if it has no  $\nu$ -atoms. A state  $\nu$  is *purely atomic*, if there is a sequence of  $\nu$ -atoms such that the sum of their probabilities is 1.

*Remark 4.8* We have  $\mathcal{A}_\varphi^\emptyset = \mathcal{A}_\varphi^\Omega = \mathcal{A}$  and  $A \in \mathcal{A}_\varphi^A \Leftrightarrow \varphi(A) \in \{0, 1\}$ . Moreover, if  $\varphi : \mathcal{A} \rightarrow [0, 1]$ , then  $\mathcal{A}_\varphi^A = \mathcal{A}$  for all  $A \in \mathcal{A}$  with  $\varphi(A) \in \{0, 1\}$ . If  $\varphi$  is nonatomic, then there exists nonempty  $A \in \mathcal{A}$  with  $\varphi(A) = 0$  [20].

**Theorem 4.9**  $\mathcal{A}_\varphi^A$  *is an asymmetric logic.*

*Proof* We show that for  $B, C \in \mathcal{A}_\varphi^A$  the following conditions are equivalent:

- (i)  $B \Delta C \in \mathcal{A}_\varphi^A$ ;
- (ii)  $B \cap C \in \mathcal{A}_\varphi^A$ .

Recall that  $\varphi(A \cap B) = \varphi(A)\varphi(B)$  and  $\varphi(A \cap C) = \varphi(A)\varphi(C)$ . The implication (i) $\Rightarrow$ (ii): we have

$$\varphi(A \cap (B \Delta C)) = \varphi(A)\varphi(B \Delta C) = \varphi(A)(\varphi(B) + \varphi(C) - 2\varphi(B \cap C)) \tag{2}$$

and via distributivity of the intersection with respect to the symmetric difference

$$\begin{aligned} \varphi(A \cap (B \Delta C)) &= \varphi((A \cap B) \Delta (A \cap C)) \\ &= \varphi(A \cap B) + \varphi(A \cap C) - 2\varphi(A \cap B \cap C) \\ &= \varphi(A)\varphi(B) + \varphi(A)\varphi(C) - 2\varphi(A \cap B \cap C). \end{aligned}$$

Now via (2) we obtain  $\varphi(A \cap (B \cap C)) = \varphi(A)\varphi(B \cap C)$ , as desired.

The implication (ii) $\Rightarrow$ (i) can be verified by inversion of the chain of arguments given above. □

**Corollary 4.10** *If a concrete logic  $\mathcal{A}_\varphi^A$  is a symmetric logic, then it is a Boolean algebra.*

**Corollary 4.11** *For  $n \geq 2$  the symmetric logic  $\mathcal{E}_{2n}^{\text{even}}$  cannot be represented in the form  $\mathcal{A}_\varphi^A$  with some  $\mathcal{A}$ ,  $\varphi$  and  $A \in \mathcal{A}$ .*

**Proposition 4.12** *Let  $\mathcal{A}$  be a Boolean algebra and  $\varphi, \psi \in P(\mathcal{A})$  be so that at least one of them is nonatomic. If  $\mathcal{A}_\varphi^A = \mathcal{A}_\psi^A$  for all  $A \in \mathcal{A}$ , then  $\varphi = \psi$ .*

*Proof* Note that  $\varphi, \psi$  have identical independent events (i.e. for any pair of events  $A$  and  $B$ ,  $\varphi(A \cap B) = \varphi(A)\varphi(B)$  if and only if  $\psi(A \cap B) = \psi(A)\psi(B)$ ) and apply Theorem 1 of [9]. □

*Example 4.13* Let  $\mathcal{A} = 2^{\Omega_6}$ ,  $\varphi(X) = \frac{1}{6} \text{card } X$  for  $X \in \mathcal{A}$ . Let  $A = \{2, 4, 6\}$ . Then

$$\mathcal{A}_\varphi^A = \{\emptyset, \Omega_6, B = \{1, 2\}, C = \{1, 4\}, D = \{1, 6\}, E = \{2, 3\}, F = \{2, 5\},$$

$$G = \{3, 4\}, H = \{3, 6\}, I = \{4, 5\}, J = \{5, 6\}, B^c, C^c, D^c, E^c, F^c, G^c, H^c, I^c, J^c\}.$$

We have  $B^c \Delta H = I$  and  $B \Delta C \notin \mathcal{A}_\varphi^A \subset \mathcal{E}_6^{\text{even}}$ .

*Example 4.14* Let  $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$ ,  $\mathcal{A} = 2^{\mathbb{N}_0}$  and a state  $\varphi$  be defined by a non-increasing sequence  $a_n = \varphi(\{n\})$ ,  $n \in \mathbb{N}_0$ . If  $a_{n+1} \leq a_n^2$  holds for all  $n \in \mathbb{N}_0$ , then there are no (nontrivial) independent events in this probability space [33, Example 1.1]. Thus  $\mathcal{A}_\varphi^A = \{\emptyset, \mathbb{N}_0\}$  for all  $A \in \mathcal{A} \setminus \{\emptyset, \mathbb{N}_0\}$ .

*Remark 4.15* The range of a purely atomic probability measure can easily be the whole  $[0, 1]$ , e.g. if the probability of the  $n$ -th atom is  $a_n = 1/2^{n+1}$ . If the range  $\{\varphi(A) : A \in \mathcal{A}\}$  of a probability measure  $\varphi$  contains the whole interval  $[0, 1]$  or at least if the range contains an arbitrary small interval  $[0, \varepsilon]$ ,  $\varepsilon > 0$ , then there are infinitely many independent events in the underlying probability space [33, Theorem 1.1].

### 4.3 When All States are $\Delta$ -subadditive

All states on Boolean algebras are subadditive and hence  $\Delta$ -subadditive.

*Problem 4.16* [6, Problem 7.1] Let  $\mathcal{E}$  be a symmetric logic such that any state  $m \in P(\mathcal{E})$  is  $\Delta$ -subadditive. Is it true that  $\mathcal{E}$  is a Boolean algebra?

A positive answer was given in [7, Theorem 4.3] with a proof by induction on the cardinality of the domain. Here we present a more general result with a new proof which is shorter and constructive—we describe the state which violates  $\Delta$ -subadditivity.

Let us recall that a state  $m_x$  on a concrete logic  $\mathcal{E}$  of subsets of  $\Omega$  is *concentrated* in a point  $x \in \Omega$  if

$$m_x(A) = \begin{cases} 1 & \text{if } x \in A, \\ 0 & \text{otherwise.} \end{cases}$$

**Theorem 4.17** *Let  $\mathcal{E}$  be a finite symmetric logic with the following property:*

*Each state on  $\mathcal{E}$  which is an affine combination of concentrated states is  $\Delta$ -subadditive.*

*Then  $\mathcal{E}$  is a Boolean algebra.*

*Proof* Suppose that  $\mathcal{E}$  is a finite symmetric logic of subsets of  $\Omega$ . Without loss of generality, we assume that  $\mathcal{E}$  satisfies

$$\forall a, b \in \Omega, a \neq b \exists A \in \mathcal{E} : a \in A, b \notin A.$$

This means that each two points  $a, b \in \Omega$  can be separated by an element of  $\mathcal{E}$ . Such a representation can be always found by the identification of points which cannot be separated. As  $\mathcal{E}$  is finite, so is  $\Omega$ . Let  $n = \text{card } \Omega$ .

For  $x \in \Omega$ , we define

$$\mathcal{E}_x = \{A \in \mathcal{E} \mid x \in A\}.$$

According to our assumptions,  $\bigcap \mathcal{E}_x = \{x\}$  for all  $x \in \Omega$ .

If  $\mathcal{E}$  contains all singletons, it is a Boolean algebra isomorphic to  $2^\Omega$ . Suppose that  $\{x\} \notin \mathcal{E}$ . We choose two sets  $A, B \in \mathcal{E}_x$  such that their intersection,  $A \cap B$ , has the least possible cardinality, say  $k$ .

*Claim*  $A \cap B$  is a proper subset of  $A$  and  $B$ , i.e., there exist  $a \in A \setminus B, b \in B \setminus A$ .

*Proof of the claim* If  $A \subset B$  and  $\text{card } A > 1$ , then there is a  $c \in A, c \neq x$ . As  $c$  can be separated from  $x$ , there is a  $C \in \mathcal{E}$  such that  $x \in C, c \notin C$ . The intersection  $A \cap C$  contains  $x$  and has a lower cardinality than  $A = A \cap B$ , a contradiction.

As a corollary, we get the following:

*Claim* Each set from  $\mathcal{E}$  has at least  $k + 1$  elements.

Now we are ready to finish the proof of the theorem. We define  $m$  as the following affine combination of concentrated states:

$$m = \frac{-k}{n - k - 1} m_x + \frac{1}{n - k - 1} \sum_{y \neq x} m_y,$$

where the sum is taken over all  $y \in \Omega \setminus \{x\}$ . Due to the preceding claim,  $m$  is non-negative. As an affine combination of states,  $m$  is additive and satisfies  $m(\Omega) = 1$ , thus it is a state. However,  $m$  is not  $\Delta$ -subadditive because

$$\begin{aligned} m(A) &= \frac{1}{n - k - 1} (-k + \text{card } A - 1), \\ m(B) &= \frac{1}{n - k - 1} (-k + \text{card } B - 1), \\ m(A) + m(B) &= \frac{1}{n - k - 1} (-2k + \text{card } A + \text{card } B - 2), \\ m(A \Delta B) &= \frac{1}{n - k - 1} (-2k + \text{card } A + \text{card } B) > m(A) + m(B). \end{aligned}$$

□

*Remark 4.18* Theorem 4.17 cannot be extended to infinite symmetric logics, see Proposition 4.8 of [7].

**Acknowledgments** The second author was supported by the Czech Technical University in Prague under project SGS12/187/OHK3/3T/13.

## References

1. Bikchentaev, A.M.: Representation of linear operators in a Hilbert space in the form of finite sums of products of projectors. *Russian Acad. Sci. Dokl. Math.* **68**(3), 376–379 (2003)
2. Bikchentaev, A.M., Sherstnev, A.N.: Projective convex combinations in  $C^*$ -algebras with the unitary factorization property. *Math. Notes* **76**(3–4), 578–581 (2004)
3. Bikchentaev, A.M.: On the representation of elements of a von Neumann algebra in the form of finite sums of products of projections. *Siberian Math. J.* **46**(1), 24–34 (2005)
4. Bikchentaev, A.M.: Projective convex combinations in  $C^*$ -algebras and the invariant subspace problem. I. *Math. Notes* **79**(1–2), 285–290 (2006)
5. Bikchentaev, A.M.: States on symmetric logics: conditional probability and independence. *Lobachevskii J. Math.* **30**(2), 101–106 (2009)
6. Bikchentaev, A.M., Yakushev, R.S.: States on symmetric logics: conditional probability and independence II. *Internat. J. Theoret. Phys.* **53**(2), 397–408 (2014)
7. Bikchentaev, A.M., Navara, M.: States on symmetric logics: extensions and metrics. *Math. Slovaca*, accepted
8. Bratteli, O., Robinson, D.W.: *Operator Algebras and Quantum Statistical Mechanics. Vol. I.* Springer, Heidelberg Berlin (1979)
9. Chen, Z., Rubin, H., Vitale, R.A.: Independence and determination of probabilities. *Proc. Amer. Math. Soc.* **125**(12), 3721–3723 (1997)
10. De Simone, A., Navara, M., Pták, P.: Extending states on finite concrete logics. *Internat. J. Theoret. Phys.* **46**(8), 2046–2052 (2007)
11. De Simone, A., Navara, M., Pták, P.: States on systems of sets that are closed under symmetric difference. Preprint (2013)
12. Flachsmeyer, J.: Note on orthocomplemented posets. In: *Proc. Conf. Topology and Measure V*, Binz, pp. 65–75. Ernst-Moritz-Arndt University, Greifswald (1982)
13. Grigoryan, S.A., Mushtari, D.H., Ovchinnikov, P.G.: Disjunctivity and alternativity in projection logics. *Internat. J. Theoret. Phys.* **39**(3), 705–709 (2000)
14. Gudder, S.: *Stochastic Methods in Quantum Mechanics.* North-Holland, New York (1979)
15. Kadison, R.V., Ringrose, J.R.: *Fundamentals of the Theory of Operator Algebras. Vol. I. Elementary theory.* Pure and Applied Mathematics, 100. Academic Press, New York (1983)
16. Kalinin, V.V.: Orthomodular partially ordered sets with dimension. (*Russian*) *Algebra and Logics* **15**(5), 535–557 (1976)
17. Kalmbach, G.: *Orthomodular Lattices.* Academic Press, London (1983)
18. Katrnoška, F.: *Logics and States of Physical Systems.* Ph.D. Thesis. Czech Technical University, Prague (1980)
19. Katrnoška, F.: Logics that are generated by idempotents. *Lobachevskii J. Math.* **15**, 11–19 (2004)
20. Marczewski, E.: Remarks on the convergence of measurable sets and measurable functions. *Colloq. Math.* **3**(2), 118–124 (1955)
21. Matoušek, M., Pták, P.: Symmetric difference on orthomodular lattices and  $Z_2$ -valued states. *Comment. Math. Univ. Carolin.* **50**(4), 535–547 (2009)
22. Mayet, R., Navara, M.: Classes of logics representable as kernels of measures. In: Pilz, G. (ed.) *Contributions to General Algebra 9*, pp. 241–248. Teubner, Stuttgart/Wien (1995)
23. Müller, V.: Jauch–Piron states on concrete quantum logics. *Internat. J. Theoret. Phys.* **32**(3), 433–442 (1993)
24. Murphy, G.J.:  *$C^*$ -algebras and Operator Theory.* Academic Press, Inc., Boston (1990)
25. Mushtari, D.H.: Logics of projectors in Banach spaces. *Soviet Math. (Iz. VUZ)* **33**(8), 59–70 (1989)
26. Mushtari, D.H.: Quantum logics of idempotents or projections in  $C^*$ -algebras. *Lobachevskii J. Math.* **33**(1), 1–4 (2012)
27. Navara, M.: Quantum logics representable as kernels of measures. *Czechoslovak Math. J.* **46**(4), 587–597 (1996)
28. Ovchinnikov, P.G., Sultanbekov, F.F.: Finite concrete logics: their structure and measures on them. *Internat. J. Theoret. Phys.* **37**(1), 147–153 (1998)

29. Ovchinnikov, P.G.: Measures on finite concrete logics. Proc. Amer. Math. Soc. **127**(7), 1957–1966 (1999)
30. Pták, P.: Some nearly Boolean orthomodular posets. Proc. Amer. Math. Soc. **126**(7), 2039–2046 (1998)
31. Pták, P.: Concrete quantum logics. Internat. J. Theoret. Phys. **39**(3), 827–837 (2000)
32. Sultanbekov, F.F.: Set logics and their representations. Internat. J. Theoret. Phys. **32**(11), 2177–2186 (1993)
33. Székely, G.J., Móri, T.F.: Independence and atoms. Proc. Amer. Math. Soc. **130**(1), 213–216 (2002)
34. Takesaki, M.: Theory of Operator Algebras, vol. I. Springer, Heidelberg – Berlin (1979)