

METRICS ON PROJECTIONS OF THE VON NEUMANN ALGEBRA ASSOCIATED WITH TRACIAL FUNCTIONALS

A. M. Bikchentaev

UDC 517.98

Abstract: Let φ be a positive functional on a von Neumann algebra \mathcal{A} and let \mathcal{A}^{pr} be the projection lattice in \mathcal{A} . Given $P, Q \in \mathcal{A}^{\text{pr}}$, put $\rho_\varphi(P, Q) = \varphi(|P - Q|)$ and $d_\varphi(P, Q) = \varphi(P \vee Q - P \wedge Q)$. Then $\rho_\varphi(P, Q) \leq d_\varphi(P, Q)$ and $\rho_\varphi(P, Q) = d_\varphi(P, Q)$ provided that $PQ = QP$. The mapping ρ_φ (or d_φ) meets the triangle inequality if and only if φ is a tracial functional. If τ is a faithful tracial functional then ρ_τ and d_τ are metrics on \mathcal{A}^{pr} . Moreover, if τ is normal then $(\mathcal{A}^{\text{pr}}, \rho_\tau)$ and $(\mathcal{A}^{\text{pr}}, d_\tau)$ are complete metric spaces. Convergences with respect to ρ_τ and d_τ are equivalent if and only if \mathcal{A} is abelian; in this case $\rho_\tau = d_\tau$. We give one more criterion for commutativity of \mathcal{A} in terms of inequalities.

DOI: 10.1134/S003744661906003X

Keywords: Hilbert space, bounded linear operator, von Neumann algebra, projection, commutativity, normal functional, state, trace

1. Definitions and Notations

Let \mathcal{H} be a Hilbert space over the field \mathbb{C} , let $\mathcal{B}(\mathcal{H})$ be the $*$ -algebra of all bounded linear operators in \mathcal{H} , let $\text{pr}(X)$ be the projection to the closure of the range of $X \in \mathcal{B}(\mathcal{H})$, and let I be the identity operator on \mathcal{H} . If $P, Q \in \mathcal{B}(\mathcal{H})^{\text{pr}}$, then $P^\perp = I - P \in \mathcal{B}(\mathcal{H})^{\text{pr}}$ and the projection $P \wedge Q$ is defined as $(P \wedge Q)\mathcal{H} = P\mathcal{H} \cap Q\mathcal{H}$, while $P \vee Q = (P^\perp \wedge Q^\perp)^\perp$ projects to $\overline{\text{lin}(P\mathcal{H} \cup Q\mathcal{H})}$. The commutant of $\mathcal{X} \subset \mathcal{B}(\mathcal{H})$ is

$$\mathcal{X}' = \{Y \in \mathcal{B}(\mathcal{H}) : XY = YX \text{ for all } X \in \mathcal{X}\}.$$

A von Neumann algebra acting on \mathcal{H} is a $*$ -subalgebra \mathcal{A} of $\mathcal{B}(\mathcal{H})$ satisfying $\mathcal{A} = \mathcal{A}''$. Given a von Neumann algebra \mathcal{A} , we denote the subset of positive elements of \mathcal{A} by \mathcal{A}^+ and the projection lattice, by \mathcal{A}^{pr} .

Given $P, Q \in \mathcal{A}^{\text{pr}}$, write $P \sim Q$ (the Murray–von Neumann equivalence) if $P = U^*U$ and $Q = UU^*$ for some $U \in \mathcal{A}$. Projections $P, Q \in \mathcal{A}$ are called *isoclinic* (we write $P \overset{\theta}{\sim} Q$ for the angle $\theta \in (0, \pi/2)$) if $PQP = \cos^2 \theta P$ and $QPQ = \cos^2 \theta Q$. If $A \in \mathcal{A}$, then $|A| = \sqrt{A^*A} \in \mathcal{A}^+$ and the projection $\text{pr}(A)$ to the closure of the range of A lies in \mathcal{A} . A positive functional φ on a von Neumann algebra \mathcal{A} is called *faithful* if $\varphi(A) = 0$ ($A \in \mathcal{A}^+$) $\Rightarrow A = 0$; while φ is *tracial* if $\varphi(Z^*Z) = \varphi(ZZ^*)$ for all $Z \in \mathcal{A}$; *normal* if $A_i \nearrow A$ ($A_i, A \in \mathcal{A}^+$) $\Rightarrow \varphi(A) = \sup_i \varphi(A_i)$; and φ is a *state* if $\varphi(I) = 1$.

2. Metrics on \mathcal{A}^{pr} Associated with a Tracial State

Lemma 1 [1, Proposition 4.4]. *The real function $\lambda \mapsto \sqrt{\lambda}$ is operator monotone on \mathbb{R}^+ .*

Let \mathcal{A} be a von Neumann algebra. Given $P, Q \in \mathcal{A}^{\text{pr}}$, put

$$P \circ Q = 2^{-1}(PQ + QP), \quad P \ominus Q = P \vee Q - P \wedge Q.$$

We have $P \circ Q \in \mathcal{A}^+ \Leftrightarrow PQ = QP$ (see the lemma in [2]). If $U \in \mathcal{A}$ is unitary, then

$$U(P \circ Q)U^* = (UPU^*) \circ (UQU^*), \quad U(P \ominus Q)U^* = (UPU^*) \ominus (UQU^*).$$

The research was funded by the subsidy allocated to Kazan Federal University for the state assignment in the sphere of scientific activities, project 1.9773.2017/8.9.

Lemma 2. Let \mathcal{A} be a von Neumann algebra and $P, Q \in \mathcal{A}^{\text{pr}}$. Then

- (i) $|P \circ Q|^2 \leq 4^{-1}(P + Q)^2$;
- (ii) $P \wedge Q \leq |P \circ Q| \leq 2^{-1}(P + Q) \leq P \vee Q$;
- (iii) $P \ominus Q = Q \ominus P = \text{pr}(|P - Q|)$;
- (iv) $P \ominus Q = P^\perp \ominus Q^\perp$;
- (v) $|P - Q|^a \leq P \ominus Q$ for all $a > 0$;
- (vi) $I \ominus P = P^\perp$, $0 \ominus P = P$, $P \ominus P^\perp = I$, and $P \ominus P = 0$;
- (vii) if $PQ = QP$, then $P \ominus Q = |P - Q|$.

PROOF. It suffices to verify the inequality $P \wedge Q \leq |P \circ Q|$ and item (i) for rank one projection $P, Q \in \mathbb{M}_2(\mathbb{C})$ (see the proof of Theorem 1 in [3] or Theorem 1 in [4]). Without loss of generality, put

$$P = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad Q = \begin{pmatrix} t & \delta\sqrt{t(1-t)} \\ \delta\sqrt{t(1-t)} & 1-t \end{pmatrix}$$

for $\delta \in \mathbb{C}$ with $|\delta| = 1$ and $0 \leq t \leq 1$. Now, the inequality $P \wedge Q \leq |P \circ Q|$ is obvious and

$$(PQ + QP)^2 = t(P + Q)^2 \leq (P + Q)^2. \quad (1)$$

By Lemma 1, from (1) we obtain $|P \circ Q| \leq 2^{-1}(P + Q) \leq P \vee Q$. Items (iii) and (v) were established in Theorem 2 of [5]. Since $P - Q = Q^\perp - P^\perp$, (iv) follows from (iii). Item (vii) is established in Proposition 1(iii) of [5]. The lemma is proven. \square

REMARK 1. Let us give a simple proof of the inequality $|P \circ Q| \leq P \vee Q$. We have $(P \pm Q)^2 \geq 0$ and $-2P \vee Q \leq -P - Q \leq PQ + QP \leq P + Q \leq 2P \vee Q$, i.e., $-P \vee Q \leq P \circ Q \leq P \vee Q$. Then $|P \circ Q| \leq P \vee Q$ by Theorem 2.4 of [6].

DEFINITION. Given a positive functional φ on a von Neumann algebra \mathcal{A} , we introduce the mappings $\rho_\varphi, d_\varphi : \mathcal{A}^{\text{pr}} \times \mathcal{A}^{\text{pr}} \rightarrow \mathbb{R}^+$ by the formulas

$$\rho_\varphi(P, Q) = \varphi(|P - Q|) \quad \text{and} \quad d_\varphi(P, Q) = \varphi(P \ominus Q) \quad \text{for all } P, Q \in \mathcal{A}^{\text{pr}}.$$

Proposition 1. The following are valid:

- (i) $\rho_\varphi(P, Q) \leq d_\varphi(P, Q)$ for all $P, Q \in \mathcal{A}^{\text{pr}}$;
- (ii) $d_\varphi(P, Q) \leq \sin^{-2} \theta \rho_\varphi(P, Q)$ for all $P, Q \in \mathcal{A}^{\text{pr}}$ with $P \overset{\theta}{\approx} Q$;
- (iii) if $PQ = QP$, then $\rho_\varphi(P, Q) = d_\varphi(P, Q)$;
- (iv) $\rho_\varphi(Q, P) = \rho_\varphi(P, Q) = \rho_\varphi(P^\perp, Q^\perp)$ for all $P, Q \in \mathcal{A}^{\text{pr}}$;
- (v) $d_\varphi(Q, P) = d_\varphi(P, Q) = d_\varphi(P^\perp, Q^\perp)$ for all $P, Q \in \mathcal{A}^{\text{pr}}$;
- (vi) $\rho_\varphi(P, 0) = d_\varphi(P, 0) = \varphi(P)$ for all $P \in \mathcal{A}^{\text{pr}}$;
- (vii) $\rho_\varphi(P, Q) = d_\varphi(P, Q) = \varphi(P + Q)$ for all $P, Q \in \mathcal{A}^{\text{pr}}$ with $PQ = 0$;
- (viii) $\varphi(|P \circ Q| - P \wedge Q) \leq d_\varphi(P, Q)$ for all $P, Q \in \mathcal{A}^{\text{pr}}$;
- (ix) $\rho_\varphi(P, Q) + \rho_\varphi(Q, R) = \rho_\varphi(P, R)$ for all $P, Q, R \in \mathcal{A}^{\text{pr}}$ with $P \leq Q \leq R$;
- (x) $d_\varphi(P, Q) + d_\varphi(Q, R) = d_\varphi(P, R)$ for all $P, Q, R \in \mathcal{A}^{\text{pr}}$ with $P \leq Q \leq R$.

PROOF. Items (i), (iii), and (iv) follow from (v), (vii), and (iv) of Lemma 2 respectively. In particular, if \mathcal{A} is abelian, then $\rho_\varphi(P, Q) = d_\varphi(P, Q)$ for all $P, Q \in \mathcal{A}^{\text{pr}}$.

Show (ii). By (iii) of Theorem 10.5 of [1] $P \wedge Q = 0$ and $P \vee Q = \sin^{-2} \theta (P - Q)^2$ for $P, Q \in \mathcal{A}^{\text{pr}}$ with $P \overset{\theta}{\approx} Q$. Since $\|P - Q\| \leq 1$ for all $P, Q \in \mathcal{A}^{\text{pr}}$; therefore, $(P - Q)^2 \leq \sqrt{(P - Q)^2} = |P - Q|$.

By Lemma 2(ii), for all $P, Q \in \mathcal{A}^{\text{pr}}$ we obtain $0 \leq |P \circ Q| - P \wedge Q \leq P \ominus Q$, which implies (viii). The proposition is proven. \square

Theorem 1. Let τ be a faithful tracial functional on a von Neumann algebra \mathcal{A} . Then ρ_τ and d_τ are metrics on \mathcal{A}^{pr} .

PROOF. Take $P, Q, R \in \mathcal{A}^{\text{pr}}$. For each pair of operators $X, Y \in \mathcal{A}$ there exist some partial isometries $U, V \in \mathcal{A}$ such that $|X + Y| \leq U|X|U^* + V|Y|V^*$ [7, Theorem 2.2]. Letting $X = P - R$ and $Y = R - Q$, we obtain the triangle inequality for ρ_τ .

From (iii) of Lemma 2 it follows that $d_\tau(P, Q) = d_\tau(Q, P)$ and $d_\tau(P, Q) = 0 \Leftrightarrow P = Q$. If $A, B \in \mathcal{A}^{\text{pr}}$, then $A \vee B - B \sim B - A \wedge B$ [8, Chapter III, Theorem 1.1.3]. Consequently,

$$\tau(A \vee B) + \tau(A \wedge B) = \tau(A) + \tau(B). \quad (2)$$

Prove the triangle inequality for d_τ ; i.e.,

$$\tau(P \vee Q) - \tau(P \wedge Q) \leq \tau(P \vee R) - \tau(P \wedge R) + \tau(R \vee Q) - \tau(R \wedge Q). \quad (3)$$

From (2) $\tau(A \wedge B) = \tau(A) + \tau(B) - \tau(A \vee B)$; therefore, (3) can be rewritten as

$$\tau(P \vee Q) \leq \tau(P \vee R) + \tau(R \vee Q) - \tau(R). \quad (4)$$

Putting $A = P \vee R$ and $B = R \vee Q$ in (2), we have

$$\tau(P \vee R) + \tau(R \vee Q) = \tau(P \vee Q \vee R) + \tau((P \vee R) \wedge (R \vee Q)). \quad (5)$$

Rewrite (4) using (5):

$$\tau(P \vee Q) \leq \tau(P \vee Q \vee R) + \tau((P \vee R) \wedge (R \vee Q)) - \tau(R). \quad (6)$$

Since $P \vee Q \vee R \geq P \vee Q$ and $(P \vee R) \wedge (R \vee Q) \geq R$; therefore, (6) is valid by monotonicity of τ on \mathcal{A}^+ . Consequently, (3) is also valid. The theorem is proven. \square

Theorem 2. *If in the conditions of Theorem 1 τ is a normal state, then $(\mathcal{A}^{\text{pr}}, \rho_\tau)$ and $(\mathcal{A}^{\text{pr}}, d_\tau)$ are complete metric spaces.*

PROOF. A von Neumann algebra \mathcal{A} possesses the topology t_τ of convergence in measure \mathcal{A} (see [9]) whose fundamental system of neighborhoods of zero is constituted by the sets

$$U(\varepsilon, \delta) = \{X \in \mathcal{A} : \exists P \in \mathcal{A}^{\text{pr}} (\|XP\| \leq \varepsilon \text{ and } \tau(P^\perp) \leq \delta)\}, \quad \varepsilon > 0, \delta > 0.$$

It is well known that $\langle \mathcal{A}, t_\tau \rangle$ is a metrizable topological $*$ -algebra. Define the L_1 -norm on \mathcal{A} by putting $\|X\|_1 = \tau(|X|)$ for all $X \in \mathcal{A}$. Let $\mathcal{A}_1 = \{X \in \mathcal{A} : \|X\|_1 \leq 1\}$.

For $(\mathcal{A}^{\text{pr}}, \rho_\tau)$, the claim follows from the $\|\cdot\|_1$ -completeness of $(\mathcal{A}_1, \|\cdot\|_1)$, the continuity of the embedding $(\mathcal{A}_1, \|\cdot\|_1)$ in the topological $*$ -algebra (\mathcal{A}, t_τ) , and the t_τ -closedness of \mathcal{A}^{pr} .

For $(\mathcal{A}^{\text{pr}}, d_\tau)$, the claim follows from coincidence of d_τ to the restriction to \mathcal{A}^{pr} of the well-known metric $d_{s,\tau}(A, B) = \tau(\text{pr}(|A - B|))$, with $A, B \in \mathcal{A}$, the continuity of the embedding $(\mathcal{A}, d_{s,\tau})$ in (\mathcal{A}, t_τ) (see [10, 11]), and the t_τ -closedness of \mathcal{A}^{pr} . The theorem is proven. \square

Theorem 3. *Let τ be a faithful tracial functional on a von Neumann algebra \mathcal{A} . Convergences with respect to the metrics ρ_τ and d_τ are equivalent if and only if \mathcal{A} is abelian.*

PROOF. If \mathcal{A} is abelian, then $\rho_\tau(P, Q) = d_\tau(P, Q)$ for all $P, Q \in \mathcal{A}^{\text{pr}}$ by assertion (iii) of Proposition 1.

If \mathcal{A} is not abelian, then \mathcal{A} has a $*$ -subalgebra $*$ -isomorphic to the complete matrix algebra $\mathbb{M}_2(\mathbb{C})$. Consider the sequence of rank one projections

$$P_n = \begin{pmatrix} \frac{1}{n} & \sqrt{\frac{1}{n} \left(1 - \frac{1}{n}\right)} \\ \sqrt{\frac{1}{n} \left(1 - \frac{1}{n}\right)} & 1 - \frac{1}{n} \end{pmatrix}, \quad n \in \mathbb{N},$$

in $\mathbb{M}_2(\mathbb{C})$. Then $P_n \rightarrow \text{diag}(0, 1)$ as $n \rightarrow \infty$ in ρ_τ , but $\{P_n\}_{n=1}^\infty$ is not fundamental in the metric d_τ , since $P_n \vee P_m = I$ and $P_n \wedge P_m = 0$ for $n \neq m$ for all $n \in \mathbb{N}$. The theorem is proven. \square

3. Characterization of Tracial Functionals

Show that ρ_φ (or d_φ) satisfies the triangle inequality if and only if φ is tracial. Let $P, Q, R \in \mathcal{A}^{\text{pr}}$. The triangle inequalities for ρ_φ and d_φ with $R = 0$ take the form $\rho_\varphi(P, Q) \leq \varphi(P+Q)$ and $d_\varphi(P, Q) \leq \varphi(P+Q)$ respectively.

Theorem 4. *For a positive normal functional φ on a von Neumann algebra \mathcal{A} the following are equivalent:*

- (i) φ is tracial;
- (ii) $\rho_\varphi(P, Q) \leq \varphi(P + Q)$ for all $P, Q \in \mathcal{A}^{\text{pr}}$;
- (iii) $d_\varphi(P, Q) \leq \varphi(P + Q)$ for all $P, Q \in \mathcal{A}^{\text{pr}}$.

PROOF. (i) \implies (ii) is established in the proof of Theorem 1.

(i) \implies (iii) follows from (2).

(ii) \implies (i) is established in item (v) of Theorem 3.4 of [12].

Below, we show that the proof of (iii) \implies (i) for an arbitrary von Neumann algebra is reduced to the case of the algebra $\mathbb{M}_2(\mathbb{C})$, in the same manner as it was done in a series of other similar cases (see [13] or [14]).

It is well known [13] that a positive normal functional φ on a von Neumann algebra \mathcal{A} is tracial if and only if $\varphi(P) = \varphi(Q)$ for all $P, Q \in \mathcal{A}^{\text{pr}}$ with $PQ = 0$ and $P \sim Q$ (see also [14, Lemma 2]). Let the $*$ -algebra \mathcal{B} in the reduced algebra $(P + Q)\mathcal{A}(P + Q)$ be generated by the partial isometry $V \in \mathcal{A}$ realizing the equivalence of P and Q . Then \mathcal{B} is $*$ -isomorphic to $\mathbb{M}_2(\mathbb{C})$ and the inequality in (ii) remains valid for the operators in \mathcal{B} and the restriction $\varphi|_{\mathcal{B}}$. Show that this restriction is a tracial functional on \mathcal{B} which implies that $\varphi(P) = \varphi(Q)$.

It is well known that each linear functional φ on $\mathbb{M}_2(\mathbb{C})$ can be presented as $\varphi(\cdot) = \text{tr}(S_\varphi \cdot)$. The matrix $S_\varphi \in \mathbb{M}_2(\mathbb{C})$ is called the *density matrix* for φ . Following the proof of Theorem 4 of [15], suppose that S_φ has two eigenvalues λ and μ , while u and v are the corresponding mutually orthogonal eigenvectors. Let P_w be an orthogonal projection to the straight line $\mathbb{C}w$ and $\varepsilon > 0$ is an arbitrary positive real. Choose linearly independent vectors x and y so that $|\varphi(P_x - P_w)| < \varepsilon$ and $|\varphi(P_y - P_w)| < \varepsilon$. Note that $P_x \vee P_y = I$ and $P_x \wedge P_y = 0$. Since

$$\lambda + \mu = \text{tr}(S_\varphi) = \varphi(I) = \varphi(P_x \vee P_y - P_x \wedge P_y) \leq \varphi(P_x) + \varphi(P_y) \leq 2\varphi(P_w) + 2\varepsilon = 2\mu + 2\varepsilon,$$

we obtain $\lambda \leq \mu + 2\varepsilon$. Since ε is arbitrary while λ and μ are interchangeable, $\lambda = \mu$. The theorem is proven. \square

For other characterizations of a trace, see [16–18] and the references therein.

REMARK 2. It is well known that $d_\varphi(P, Q) \leq \varphi(P + Q)$ for every positive normal functional φ on a von Neumann algebra \mathcal{A} and every pair of commutative projections $P, Q \in \mathcal{A}^{\text{pr}}$ [19, p. 168]. By (iii) of Proposition 1, $\rho_\varphi(P, Q) \leq \varphi(P + Q)$ for every positive normal functional φ on a von Neumann algebra \mathcal{A} and each pair of commutative projections $P, Q \in \mathcal{A}^{\text{pr}}$. Theorem 4 demonstrates that each of these inequalities holds for all pairs $P, Q \in \mathcal{A}^{\text{pr}}$ if and only if φ is tracial.

Corollary. *For a von Neumann algebra \mathcal{A} the following are equivalent:*

- (i) \mathcal{A} is abelian;
- (ii) $\rho_\varphi(P, Q) \leq \varphi(P + Q)$ for all normal states φ on \mathcal{A} and $P, Q \in \mathcal{A}^{\text{pr}}$;
- (iii) $d_\varphi(P, Q) \leq \varphi(P + Q)$ for all normal states φ on \mathcal{A} and $P, Q \in \mathcal{A}^{\text{pr}}$.

PROOF. By Theorem 4, every normal state on \mathcal{A} is tracial. The set of normal states on \mathcal{A} separates the points of \mathcal{A} [8, Chapter III, Theorem 2.4.5]; therefore, \mathcal{A} is commutative. \square

The author is grateful to the referee for valuable advice.

References

1. Sherstnev A. N., *Methods of Bilinear Forms in Noncommutative Measure and Integral Theory* [Russian], Fizmatlit, Moscow (2008).
2. Uchiyama M., “Commutativity of selfadjoint operators,” *Pacific J. Math.*, vol. 161, no. 2, 385–392 (1993).
3. Bikchentaev A. M., “Commutation of projections and trace characterization on von Neumann algebras. II,” *Math. Notes*, vol. 89, no. 4, 461–471 (2011).
4. Bikchentaev A. M., “Commutativity of operators and characterization of traces on C^* -algebras,” *Dokl. Math.*, vol. 87, no. 1, 79–82 (2013).
5. Bikchentaev A. M., “Differences of idempotents in C^* -algebras,” *Sib. Math. J.*, vol. 58, no. 2, 183–189 (2017).
6. Bikchentaev A. M., “On hermitian operators X and Y meeting the condition $-Y \leq X \leq Y$,” *Lobachevskii J. Math.*, vol. 34, no. 3, 227–233 (2013).
7. Akemann C. A., Anderson J., and Pedersen G. K., “Triangle inequalities in operator algebras,” *Linear Multilinear Algebra*, vol. 11, no. 2, 167–178 (1982).
8. Blackadar B., *Operator Algebras. Theory of C^* -Algebras and von Neumann Algebras. Operator Algebras and Non-Commutative Geometry. III*, Springer-Verlag, Berlin (2006) (Encycl. Math. Sci.; V. 122).
9. Nelson E., “Notes on non-commutative integration,” *J. Funct. Anal.*, vol. 15, no. 2, 103–116 (1974).
10. Ciach L. J., “Linear-topological spaces of operators affiliated with a von Neumann algebra,” *Bull. Acad. Pol. Sci. Math.*, vol. 31, no. 3–4, 161–166 (1983).
11. Bikchentaev A. M., “Minimality of convergence in measure topologies on finite von Neumann algebras,” *Math. Notes*, vol. 75, no. 3, 315–321 (2004).
12. Bikchentaev A. M., “Commutativity of projections and characterization of traces on von Neumann algebras,” *Sib. Math. J.*, vol. 51, no. 6, 971–977 (2010).
13. Gardner L. T., “An inequality characterizes the trace,” *Canad. J. Math.*, vol. 31, no. 6, 1322–1328 (1979).
14. Tikhonov O. E., “Subadditivity inequalities in von Neumann algebras and characterization of tracial functionals,” *Positivity*, vol. 9, no. 2, 259–264 (2005).
15. Petz D. and Zemánek J., “Characterizations of the trace,” *Linear Algebra Appl.*, vol. 111, 43–52 (1988).
16. Bikchentaev A. M., “Commutation of projections and characterization of traces on von Neumann algebras. III,” *Int. J. Theor. Phys.*, vol. 54, no. 12, 4482–4493 (2015).
17. Bikchentaev A. M., “Inequality for a trace on a unital C^* -algebra,” *Math. Notes*, vol. 99, no. 4, 487–491 (2016).
18. Bikchentaev A. M., “Trace and differences of idempotents in C^* -algebras,” *Math. Notes*, vol. 105, no. 5, 641–648 (2019).
19. Kadison R. V. and Ringrose J. R., *Fundamentals of the Theory of Operator Algebras. V. I. Elementary Theory*, Acad. Press, New York and London (1983) (Pure Appl. Math.; V. 100).

A. M. BIKCHENTAEV

LOBACHEVSKII INSTITUTE OF MATHEMATICS AND MECHANICS
OF KAZAN (VOLGA REGION) FEDERAL UNIVERSITY, KAZAN, RUSSIA

E-mail address: Airat.Bikchentaev@kpfu.ru